

# Assessing the potential of wildfires as a sustainable bioenergy opportunity

SANTIAGO R. VERÓN\*, ESTEBAN G. JOBBÁGY†, CARLOS M. DI BELLA\*, JOSÉ M. PARUELO‡ and ROBERT B. JACKSON§

\*Instituto de Clima y Agua, INTA-CONICET, Castelar, 1712, Argentina, †Grupo de Estudios Ambientales-IMASL, Universidad Nacional de San Luis-CONICET, Avenida Ejército de los Andes 950, San Luis, 5700, Argentina, ‡Laboratorio de Análisis Regional y Teledetección, Facultad de Agronomía, UBA-CONICET, Buenos Aires, 1417, Argentina, §Department of Biology, Nicholas School of the Environment, and Center on Global Change-Duke University, Durham, NC, 27708-0338, USA

## Abstract

As the environmental and economic consequences of fossil-fuel use become clear, land is increasingly targeted as a source of bioenergy. We explore the potential for generating electricity from biomass vulnerable to fires as an ecologic and socioeconomic opportunity that can reduce the risk of greenhouse gas generation from wildfires and help to create incentives to preserve natural and seminatural vegetation and prevent its conversion to agriculture, including biofuel crops. On the basis of a global analysis of the energy generation and spatial distribution of fires, we show that between 2003 and 2010, global fires consumed  $\sim 8300 \pm 592$  PJ  $\text{yr}^{-1}$  of energy, equivalent to  $\sim 36$ – $44\%$  of the global electricity consumption in 2008 and  $>100\%$  national consumption in 57 countries. Forests/woodlands, cultivated areas, shrublands, and grasslands contributed 53%, 19%, 16%, and 3.5% of the global energy released by fires. Although many agroecological, socioeconomic, and engineering challenges need to be overcome before diverting the energy lost in fires into more useable forms, done cautiously it could reconcile habitat preservation with economic yields in natural systems.

**Keywords:** bioelectricity, biomass burning, dry forests, fire radiative energy

Received 19 December 2011 and accepted 24 March 2012

## Introduction

In pace with declining fossil-fuel reserves and increasing concerns on global environmental deterioration, humans are trying to diversify and expand their sources of energy. Being the predominant energy source in pre-industrial times, land ecosystems are gaining renewed attention as energy providers (Campbell *et al.*, 2009; Ohlrogge *et al.*, 2009; Richter *et al.*, 2009) at a challenging time in which the appropriation of food, fiber, and timber is already accounting for one quarter of their global net primary production (NPP) (Haberl *et al.*, 2007). Although biofuel crops receive increasing attention and government support, their expansion may compromise food production and provide questionable climate benefits (Fargione *et al.*, 2008; Searchinger *et al.*, 2008; Piñeiro *et al.*, 2009). New perspectives on which ecosystems should be targeted and how they should be managed to limit the trade-offs among energy generation, food production, and environmental protection are needed. Wildfires are an increasingly important pathway of

energy release from land ecosystems, and here, we explore the magnitude and distribution of this energy flux and discuss its potential diversion to electricity generation and some of the associated environmental benefits and costs.

Globally, between 3.3 and 4.3 million  $\text{km}^{-2}$  of land, roughly the size of India, burn each year (Giglio *et al.*, 2010). These fires combust plant biomass and release particulate matter and greenhouse gasses, including the equivalent to 2 Pg C  $\text{yr}^{-1}$  of  $\text{CO}_2$ ,  $\text{CH}_4$ , and  $\text{N}_2\text{O}$ , to the atmosphere (van der Werf *et al.*, 2010). They also disturb ecosystems and the services that they provide to society by triggering soil erosion, altering the hydrologic cycle, and, sometimes, destroying infrastructure (DeFries *et al.*, 2002; Hoffmann *et al.*, 2002, 2003; Mouillot & Field, 2005; Lohman *et al.*, 2007; van der Werf *et al.*, 2008). Throughout the history of life on Earth, fires have played a key role shaping plant adaptations, ecosystem composition and distribution (Bond *et al.*, 2005), and global biogeochemical cycles (Bergman *et al.*, 2004). Nowadays, however, in a world in which 75% of the ice-free land ecosystems are subjected to some type of management and use (Ellis & Ramankutty, 2008), their significance is very different. Wildfires today have

Correspondence: Esteban G. Jobbágy, tel. + 54 266 448 2903, fax + 54 266 448 2903, e-mail: jobbagy@unsl.edu.ar

increased their intensity, extent, and duration over many areas, favored by climate change and human activity (Kasischke *et al.*, 1995; Skinner *et al.*, 2006; West-erling *et al.*, 2006, 2011; Tymstra *et al.*, 2007; Flannigan *et al.*, 2009; Liu *et al.*, 2009; Le Page *et al.*, 2010; Pechony & Shindell, 2010; Wotton *et al.*, 2010). As a consequence, it is very likely that they have exceeded their 'natural' role as an ecologic and biogeochemical agent, becoming a new force of change in systems where they were uncommon before.

We propose that the controlled combustion for energy production, based on the diversion of biomass that would eventually burn in wildfires, provides an immediate opportunity to reduce fossil-fuel use and its associated greenhouse gas (GHG) emissions and satisfy the growing energy needs of some developing economies. The climatic benefits of this intervention not only involve replacing fossil fuels but also reducing other non-CO<sub>2</sub> gasses with high global warming potential, particularly carbon monoxide, and black carbon emissions, which are significant in wildfires, but avoidable under controlled combustion conditions (Crutzen & Andreae, 1990; Bond *et al.*, 2005; Ramanathan *et al.*, 2008). Furthermore, along with reducing net GHG emissions, biomass harvesting can provide economic returns that, ideally, can help maintain habitats vulnerable to agricultural expansion.

To explore the potential of this bioenergy alternative, we first quantify and map global energy release by wildfires and the corresponding fraction of the average NPP that burns in these fires. Next, we characterize wildfire energy release across biomes and identify countries where its partial allocation to electricity generation could satisfy a substantial portion of their electricity demand.

## Materials and methods

### Fire radiative energy (FRE)

We estimated global radiative energy released by fires from January 2003 to December 2010 using the MOD and MYD14CMG fire products (Giglio *et al.*, 2006) generated from the MODIS sensor collection 5 onboard Terra and Aqua platforms. This dataset integrates subdaily, 1 km<sup>-2</sup> resolution data into monthly values for 0.5° × 0.5° grid cells. Using a probabilistic characterization of fire density that avoids assumptions about fire duration, we calculated the FRE released (FRE, in MJ yr<sup>-1</sup>) from any given cell as follows:

$$\text{FRE}_{y,\text{cell}} = \sum_{i=1}^{i=12} \frac{\text{Raw}_i}{\text{Total}_i - \text{Cloud}_i} \times \text{sFRP}_i \times A_{\text{cell}}$$

where *i* stands for month, Raw indicates the number 1 km<sup>-2</sup> fire pixels observed by Aqua and Terra for that particular cell

and month (pixels cell<sup>-1</sup> mo<sup>-1</sup>), Total shows the number of pixels that were screened within a given cell during the month *i* (pixels cell<sup>-1</sup> mo<sup>-1</sup>), and Cloud represents the number of cloudy pixels screened in the month *i* (Aqua + Terra) (pixels cell<sup>-1</sup> mo<sup>-1</sup>). The first three terms of the equation yield a dimensionless index of fire density ranging from 0 to 1 for each cell and month. The term sFRP<sub>*i*</sub> (specific radiative power per fire pixel in MW pixel<sup>-1</sup>) corresponds to the fire radiative power values provided by the CMG dataset, which were estimated as the sum of the power release measured in all fire pixels' FRP divided by the number of fire pixels screened during a calendar month. The sFRP<sub>*i*</sub> values (MW pixel<sup>-1</sup>) were transformed to energy per month (MJ mo<sup>-1</sup> pixel<sup>-1</sup>), and final values were scaled to the whole cell by considering the number of pixels per cell (A).

Sources of uncertainty in our FRE calculation would stem from biases in the estimates of the real number of fire pixels – due to limited sampling resulting from satellite orbits, cloud masking, and limitations in instruments and detection algorithms (Giglio *et al.*, 2006) – and their average radiative power. For example, recent studies showed that improper background characterization may hamper cool fire detection (Schroeder *et al.*, 2010), and sampling artifacts (i.e., 'Bow tie effect', Freeborn *et al.*, 2011) may provide flawed FRP estimates. Our use of discrete MODIS observations to describe a continuous process assumed that the effectively observed pixels offer a reasonable representation of all pixels within any given cell and for any particular month. Similarly, we assumed that the sFRP values of the observed fire pixels were representative of those fire pixels that could have been obscured by clouds or missed by the satellite passes. To test the validity of our assumptions, we validated our methodology against the Global Fire Emissions Database v3 (van der Werf *et al.*, 2010).

### Total energy release and burned biomass

We assumed a ratio of total-to-radiative energy release from wildfires of 8.1 (radiative fraction = 12.3%), based on previously reported values of 18 MJ of total energy and 2.21 MJ of radiative energy per kg of biomass (Freeborn *et al.*, 2007; Campbell *et al.*, 2009). Nonradiative energy losses are attributable to conduction, convection, and vaporization, and other secondary processes (Wooster *et al.*, 2005). We chose radiative energy rates of 2.21 MJ kg<sup>-1</sup> (Freeborn *et al.*, 2007) as opposed to alternative figures of 2.71 MJ kg<sup>-1</sup> (Wooster *et al.*, 2005) because the former has been estimated for a wider variety of biomass types, including grasses, branches, twigs, and woody vegetation. Burned biomass was then calculated as the ratio between the total energy released by fires and the heating value of biomass (18 MJ kg<sup>-1</sup>, Campbell *et al.*, 2009). Average global NPP values were obtained from Imhoff *et al.* (2004) dataset. These authors used the CASA (Potter *et al.*, 1993) carbon model driven by a global satellite-derived vegetation index (AVHRR-NDVI) and climate data obtained between 1982 and 1998 (Imhoff *et al.*, 2004). CASA estimates NPP as a product of time-varying surface solar irradiance, NDVI computed from AVHRR sensor, a constant maximum light use efficiency modi-

fied by time-varying stress scalar terms for temperature and moisture effects. In turn, these stress scalar terms are calculated from the difference to optimal temperatures or from water deficits considering a land cover map produced by Hansen *et al.* (2000) and climatic and soil texture data taken from the International Satellite Land Surface Climatology Project, Initiative II (ISLSCP II, Hall *et al.*, 2005), and Zobler (1986), respectively. Therefore, fires or any other disturbance that reduce radiation interception by plants should be captured by this modeling approach.

### *Spatial distribution of energy consumption and agricultural fires estimation*

We assessed the spatial distribution of energy consumption globally as the product of population density and per capita energy consumption for each country. National population density data were obtained from the Gridded Population of the World Version 3 (GPWv3) (Center for International Earth Science Information Network (CIESIN), Columbia University; & Centro Internacional de Agricultura Tropical (CIAT), 2005), and the total energy consumption per capita and by country were calculated from data published by the International Energy Agency (IEA) Statistics Division (2007). To estimate the amount of fires associated with agriculture, we overlapped the FRE map with the Global Land Cover (Arino *et al.*, 2008) vegetation classes aggregated to two classes (agricultural vs. other).

## Results

The radiative energy released by fires from 2003 to 2010 averaged  $8300 \pm 562$  PJ per year, combusting  $1.8 \pm 0.1$  Pg C yr<sup>-1</sup> dry mass or 3.2% of global annual NPP (Fig. 1, see Supplementary Table 1). Africa and South America, with only 20% and 12% of total land surface area, were responsible for 48% and 24% of this energy release, respectively. In several areas of the globe – for instance sub-Saharan and tropical Africa, northern Australia, southern Russia, Kazakhstan, and southeast Asia – the amount of NPP burned in fires exceeds 20% and can be as much as 60% (Fig. 1b).

Fires in forests/woodlands/savannas, cultivated areas, shrublands, and grasslands contributed 53%, 19%, 16%, and 3.5% of the global energy release, consuming on average 2.4%, 1.9%, 4.8%, and 1.7% of NPP in those systems. The low fraction of grassland NPP consumed by fires may be due to their higher below-ground allocation (Gower *et al.*, 1999; Saugier *et al.*, 2001; Chapin *et al.*, 2002), higher biomass turnover rates (e.g. Cebrian, 1999), and herbivore consumption (McNaughton *et al.*, 1989) as compared with forested ecosystems. Approximately 81% of the global fire energy release comes from areas dominated by natural vegetation, as suggested by land cover maps (Arino *et al.*, 2008). The amounts differed markedly by country,


however; Australia and the Democratic Republic of the Congo had 90% and 93% of their fires under natural and seminatural vegetation, whereas >40% of the energy released by fires in India and China came from cultivated land. The human context left its imprint on fire energy release patterns, as suggested by contrasts within the same biome across political borders, such as those between Russia and China or Portugal and Spain (Fig. 1b).

Between 2003 and 2010, the average total energy annually released by fires equaled 14% of the total energy consumed by humans in 2008. In principle, if all of the biomass that fed these fires could be diverted to energy generation assuming an efficiency ranging from 33% to 40% conversion (efficiencies from the standard steam-Rankine cycle and conservative commercially mature power generation facilities, respectively, Jin *et al.*, 2009; Schiermeier *et al.*, 2008), the burned biomass would supply between 36% and 44% of the global electricity consumption in 2008 (Energy Information Administration, 2008).

Although many biophysical, socioeconomic, and technological factors limit the full capture of this bioenergy source, regions of greater opportunity can be identified. One important component is the geographic match between fire energy release and electricity demand (Fig. 1c), which can be assessed nationally. Among the top 12 countries of highest fire energy release, which accounted for 66% of the global total (Fig. 2), we identified three groups of countries that display contrasting ratios of this potential energy source to electricity consumption. In the first group of countries with ratios <0.2 (e.g., the United States), electricity generation with biomass could be locally important, but is unlikely to alter the national energy mix. In the second group (e.g., Argentina, Brazil, Australia, and Russia), ratios between 0.2 and 5 suggest a better opportunity to replace fossil fuels. Finally, countries with a ratio >5, such as most of the sub-Saharan African nations, have low electricity demand overall; in these countries, the bioenergy strategy that we describe may be most helpful in supplying energy for future economic growth.

## Discussion


According to our analysis, fires consume ~3.2% of terrestrial NPP, producing considerable GHG emissions and generating ~8300 PJ of biomass energy release each year. Although the coarse vegetation classes that we used do not allow a detailed quantification of the contribution of, for example, C4 grasses, at a global scale our values are consistent with, but slightly more conservative than, previously reported estimates in the literature. Our estimate of 1.8 Pg C of biomass burned annually is



**Fig. 1** Global distribution of wildfires and human energy consumption: (a) Radiative energy released by fires shown as the average energy flux ( $\text{TJ yr}^{-1}$ ) per grid cell ( $0.5$  by  $0.5$  degrees) estimated from the MOD14CMG product for the 2003–2010 period. (b) Percentage of mean net primary production (NPP) that burns in each grid cell to explain observed fire energy release rates (NPP values were obtained from the CASA model (Imhoff *et al.*, 2004)). (c) Energy use by humans ( $\text{TJ yr}^{-1}$ ) per grid cell ( $0.5$  by  $0.5$  degrees) based on the Gridded Population of the World (GPWv3) dataset and country-level estimates of per capita energy consumption from the International Energy Agency.

within the range of values reported as  $2 \text{ Pg C yr}^{-1}$  by van der Werf *et al.* (2010) and  $1.1 \text{ Pg C yr}^{-1}$  by Haberl *et al.* (2007). We validated our estimates against the most recent and comprehensive fire database (Global Fire Emissions Database v.3), wherein fire emissions are

derived from the combination of a biogeochemical model and remote sensing estimates of burned area, to estimate burnt biomass (van der Werf *et al.*, 2010). We found a very good agreement (linear  $r^2 = 0.9$ , slope = 1.1, and y-intercept =  $-8$ ) for 14 world regions and



**Fig. 2** Potential electricity supply through wildfire diversion vs. current electricity demand across nations. Estimates are based on fire energy release from agricultural (red) and non-agricultural fires (green) and current electricity consumption (black), assuming a conversion efficiency of 40%. Countries are sorted from highest to lowest fire energy released. Listed countries account for 66% of the total potential electricity. For 57 countries, this bioenergy source supply exceeds national electricity consumption.

7 years (2003–2009). In addition, we calculated the correlation coefficient between our FRE estimate and the GFEDv.3 C emissions at a per pixel ( $0.5^\circ \times 0.5^\circ$ ) basis for the 2003–2009 period, showing a correlation coefficient  $>0.6$  in more than 60% of the pixels, suggesting that our simpler approach successfully captured spatial and temporal fire energy release patterns.

The land use context of wildfires, their long-term history, and their current geographic match with human energy consumption need to be considered in the discussion of electricity generation with biomass and its environmental and social opportunities. However, it is important to consider that the disruption of fire regimes on natural systems could have contrasting effects depending on their fire history, threatening those that have been shaped by long-term recurrent fires, but protecting those that suffer increasing burning in response to recent climate change or intensifying human ignition (Kasischke *et al.*, 1995; Skinner *et al.*, 2006; Westerling

*et al.*, 2006, 2011; Tymstra *et al.*, 2007; Flannigan *et al.*, 2009; Liu *et al.*, 2009; Pausas & Keely, 2009; Pechony & Shindell, 2010; Wotton *et al.*, 2010).

A sustained and sustainable diversion of biomass from wildfires to electricity generation would require a careful categorization of fires capable of separating those that periodically affect (semi)natural vegetation (i.e. regime fires), from those that represent a one-time event associated with the clearing and replacement of (semi)natural vegetation by crops (i.e. clearing fires), and those involving the reduction in crop residues (i.e. agricultural fires). We here focused on ‘regime fires’ which may be the most attractive option for biomass diversion to electricity generation given that ‘clearing fires’ would offer a single event of biomass availability, in many cases undesirable for additional environmental reasons. ‘Agricultural fires’, in turn, demand a more detailed analysis because of the competing uses of the crop residues that they consume, which, besides bioenergy, include animal feed soil fertility maintenance through erosion control and soil organic carbon generation (Smil, 2003). According to one of the most recent global land use maps (Arino *et al.*, 2008) and our own fire distribution maps, natural and seminatural systems generated 81% of the global energy release by fires in the study period.

The geographic patterns of energy release by fires and electricity consumption by humans (Figs 1c and 2) reveal an overlap in some moderately developed countries, such as Brazil, Argentina, Russia, and Mexico, in which labor availability and harvesting costs could eventually make electricity generation with biomass feasible (Central Intelligence Agency, 2009). Although the most immediate opportunities for electricity generation using fire-prone biomass occur in these countries, Africa shows the largest potential. In the northern and southern fire belts of this continent (Fig. 1a), most countries currently have low electricity generation and rely on biomass to cover  $>75\%$  of their national energy consumption (Malimbwi & Zahabu, 2010). High urbanization rates in these countries have raised charcoal demands from an increasingly broader radius of rural areas (e.g.  $>400$  km around Dakar and Dar es Salaam) (Luoga, 2000; Malimbwi & Zahabu, 2010), making it a dominant source of rural income (Chidumayo *et al.*, 2001) and highlighting the already significant role of natural ecosystems as a major energy source in these developing economies.

Although significant and promising as a potential energy source, the massive use of fire-prone biomass from natural ecosystems introduces multiple technological and socioeconomic challenges, as well as likely trade-offs with other ecosystem services. The alteration of nutrient cycles brought by the replacement of *in situ*

biomass burning with harvesting, removal, and *ex situ* combustion could lead to net nutrient losses and deterioration of soil fertility. This is particularly feasible for nonvolatile elements such as phosphorus and base cations, which would otherwise have remained in ashes after fires, their net removal could limit productivity in highly weathered or sandy soils. Harvesting schemes that leave nutrient-rich components such as leaves and bark in place or, logistically more challenging, return ashes back from power plants to the field could reduce this potential impact. Potential long-term change in biodiversity and pollinator abundance with reduced fire frequency is another front that needs further ecological study. Evidences from the large body of literature evaluating the effects of coarse woody debris extraction (a biomass component that will likely be eliminated under the management schemes proposed here) suggest mixed effects upon ecosystem services (e.g. MacNally *et al.*, 2002; Owens *et al.*, 2008). A recent study from an arid ecosystem concluded that coarse woody debris removal only showed transient negative effects on flower visitor abundance, whereas positive or nonsignificant effects were found for seed production of the dominant tree species, cover, richness, and composition of understory plants or soil properties (Vázquez *et al.*, 2012). Future research should also elucidate how electricity generation with native vegetation biomass could best be implemented to maximize the provision of other ecosystem goods and services for local people and societies and what mechanisms could warrant sustainable harvesting schemes.

Some challenging logistical and political issues emerge with the proposed energy generation approach. The distance of fire-prone areas to centers of energy consumption (Figs 1a and 1c) could impose constraints on the feasibility of large electricity generation plants. The location (distance, accessibility), type (woody vs. herbaceous, large vs. small wood), timing (harvest-regeneration periods, biomass moisture cycles), and density (harvestable mass per unit of area) of fire-prone biomass availability will determine the feasibility, design, and labor/equipment needs (Mahmoudi *et al.*, 2009). Energy consumption and road and electricity grid networks will define the type of generation method (Kumar *et al.*, 2003; Yagi & Nakata, 2011). Although here we propose capturing only the fraction of primary productivity that would otherwise be consumed by wildfires, it would be difficult to define the exact magnitude of that fraction and tempting to exceed it once the energy generation economy starts. Although suggesting a different perspective on bioenergy opportunities in land ecosystems, our proposal does not tackle the complex issue of environmental impacts and the development of management and regulation practices that

could minimize them. That challenging task will require the application of ecologic, agronomic, and social knowledge under very different geographic contexts. Our increasing understanding of fire ecology, including recent fire prediction models (Preisler *et al.*, 2011), will aid in this process, yet it would be ultimately defined by the interaction of land managers, energy markets, and policy makers among other key stakeholders.

### Acknowledgements

This article was supported by the Inter American Institute for Global Change Research (CRN 2031, NSF Grant GEO-0452325), IDRC (International Development Research Centre of Canada), CYTED (SERENA), and the U.S. National Science Foundation (DEB 0717191). We thank Juan Viciana, Carlos Turc, and Isaac and Sabrina Buchovsky for providing information and insights. Coni Caride and German Baldi helped with image processing. SRV thanks NIN, Pixies, and PJ Harvey. We also thank Elliot Campbell, Holly Gibbs, Louis Giglio, and Prasad Kasibhatla for comments on previous versions of the manuscript.

### References

- Arino O, Bicheron P, Achard F, Latham J, Witt R, Weber JL (2008) GLOBCOVER: the most detailed portrait of Earth. *European Space Agency*, **136**, 24–31.
- Bergman NM, Lenton TM, Watson AJ (2004) COPSE: a new model of biogeochemical cycling over phanerozoic time. *American Journal of Science*, **304**, 397–437.
- Bond WJ, Woodward FI, Midgley GF (2005) The global distribution of ecosystems in a world without fire. *New Phytologist*, **165**, 525–538.
- Campbell JE, Lobell DB, Field CB (2009) Greater transportation energy and GHG offsets from bioelectricity than ethanol. *Science*, **324**, 1055–1057.
- Cebrian J (1999) Patterns in the fate of production in plant communities. *American Naturalist*, **154**, 449–468.
- Center for International Earth Science Information Network (CIESIN), Columbia University; and Centro Internacional de Agricultura Tropical (CIAT) (2005) Gridded Population of the World Version 3 (GPWv3). Socioeconomic Data and Applications Center (SEDAC), Columbia University, Palisades, NY. Available at: <http://sedac.ciesin.columbia.edu/gpw> (accessed 23 January 2009).
- Central Intelligence Agency (2009) *The World Factbook*. CIA, Langley, VA. Available at: <https://www.cia.gov/library/publications/the-world-factbook/> (accessed 19 January 2010).
- Chapin FSJ, Matson PA, Mooney HA (2002) *Principles of Terrestrial Ecology*. Springer-Verlag, New York, NY, USA.
- Chidumayo EN, Masaiteti I, Ntalasha H, Kalumiana OS (2001) *Charcoal Potential in Southern Africa (CHAPOSA)*. Stockholm Environment Institute, Stockholm.
- Crutzen PJ, Andreae MO (1990) Biomass burning in the tropics: impact on atmospheric chemistry and the biogeochemical cycles. *Science*, **250**, 1669–1678.
- DeFries RS, Houghton RA, Hansen MC, Field CB, Skole D, Townshend J (2002) Carbon emissions from tropical deforestation and regrowth based on satellite observations for the 1980s and 1990s. *Proceedings of the National Academy of Sciences*, **99**, 14256–14261.
- Ellis EC, Ramankutty N (2008) Putting people in the map: anthropogenic biomes of the world. *Frontiers in Ecology and Environment*, **6**, 439–447.
- Energy Information Administration (2008) International Energy Annual 2008 Available at: <http://www.eia.doe.gov> (accessed 23 May 2011).
- Fargione J, Hill J, Tilman D, Polasky S, Hawthorne P (2008) Land clearing and the biofuel carbon debt. *Science*, **219**, 1235–1238.
- Flannigan MD, Krawchuck MA, de Groot WJ, Wotton BM, Gowman LM (2009) Implications of changing climate for global wildland fire. *International Journal of Wildland Fire*, **18**, 483–507.
- Freeborn PH, Wooster MJ, Hao WM, Ryan CA, Nordgren BL, Baker SP, Ichoku C (2007) Relationships between energy release, fuel mass loss, and trace gas and

- aerosol emissions during laboratory biomass fires. *Journal of Geophysical Research*, **113**, D01301.
- Freeborn PH, Wooster MJ, Roberts G (2011) Addressing the spatiotemporal sampling design of MODIS to provide estimates of the fire radiative energy emitted from Africa. *Remote Sensing of Environment*, **115**, 475–489.
- Giglio L, Cizar I, Justice CO (2006) Global distribution and seasonality of active fires as observed with the Terra and Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) sensors. *Journal of Geophysical Research*, **111**, JG000142.
- Giglio L, Randerson JT, van der Werf GR, Kasibhatla PS, Collatz GJ, Morton DC, DeFries RS (2010) Assessing variability and long-term trends in burned area by merging multiple satellite fire products. *Biogeosciences*, **7**, 1171–1186. doi: 10.5194/bg-7-1171-2010.
- Gower ST, Kucharik CJ, Norman JM (1999) Direct and indirect estimation of leaf area index, fAPAR, and net primary Production of terrestrial ecosystems. *Remote Sensing of the Environment*, **70**, 29–51.
- Haberl H, Erb KH, Krausmann F *et al.* (2007) Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems. *Proceedings of the National Academy of Sciences*, **104**, 12942–12947.
- Hall FG, Collatz G, Los S, Brown de Colstoun E, Landis D (Eds) (2005) ISLSCP Initiative II [DVD/CD-ROM], NASA, Washington, DC
- Hansen M, DeFries R, Townshend JRG, Sohlberg R (2000) Global land cover classification at 1 km spatial resolution using a classification tree approach. *International Journal of Remote Sensing*, **21**, 1331–1364.
- Hoffmann W, Schroeder W, Jackson RB (2002) Positive feedbacks of fire, climate, and vegetation and the conversion of tropical savanna. *Geophysical Research Letters*, **29**, 2052. doi: 10.1029/2002GL015424.
- Hoffmann W, Schroeder W, Jackson RB (2003) Regional feedbacks among fire, climate, and tropical deforestation. *Journal of Geophysical Research*, **108**, D234721.
- Imhoff ML, Bounoua L, Ricketts T, Loucks C, Harris R, Lawrence WT (2004) Global patterns in human consumption of net primary production. *Nature*, **429**, 870–873.
- International Energy Agency (IEA) Statistics Division. (2007) Energy balances of OECD countries (2008 edition) and energy balances of non-OECD countries (2007 edition). IEA, Paris. Available at: <http://data.iaea.org/ieastore/default.asp> (accessed 5 October 2009).
- Jin H, Larson ED, Celik FD (2009) Performance and cost analysis of future, commercially mature gasification-based electric power generation from switchgrass. *Biofuels, Bioproducts and Biorefining*, **3**, 142–173.
- Kasischke ES, Christinasen NL Jr, Stocks BJ (1995) Fire, global warming, and the carbon balance of boreal forests. *Ecological Applications*, **5**, 437–451.
- Kumar A, Cameron JB, Flynn PC (2003) Biomass power cost and optimum plant size in western Canada. *Biomass and Bioenergy*, **24**, 445–464.
- Le Page Y, Oom D, Silva JMN, Jönsson P, Pereira JMC (2010) Seasonality of vegetation fires as modified by human action: observing the deviation from eco-climatic fire regimes. *Global Ecology and Biogeography*, **19**, 575–588.
- Liu YQ, Stanturf J, Goodrick S (2009) Trends in global wildfire potential in a changing climate. *Forest Ecology and Management*, **259**, 685–697.
- Lohman DJ, Bickford D, Sodhi NS (2007) The burning issue. *Science*, **316**, 376.
- Luoga EJ (2000) The effect of human disturbances on diversity and dynamics of Eastern Tanzania Miombo arborescent species. Unpublished PhD thesis University of Witwatersrand, Johannesburg.
- MacNally R, Horrocks G, Pettifer L (2002) Experimental evidence for potential beneficial effects of fallen timber in forests. *Ecological Applications*, **12**, 1588–1594.
- Mahmoudi M, Sowlati T, Sokhansanj S (2009) Logistics of supplying biomass from a mountain pine beetle-infested forest to a power plant in British Columbia. *Scandinavian Journal of Forest Research*, **24**, 76–86.
- Malimbwi RE, Zahabu E (2010) REDD experience in Tanzania. In: *REDD, Forest Governance and Rural Livelihoods: The Emerging Agenda* (eds Springate-Baginski O, Wollenberg E), pp. 109–134. CIFOR, Bogor, Indonesia.
- McNaughton SJ, Oesterheld M, Frank DA, Williams KJ (1989) Ecosystem-level patterns of primary productivity and herbivory in terrestrial habitats. *Nature*, **341**, 142–144.
- Mouillot F, Field CB (2005) Fire history and the global carbon budget: a 1° × 1° fire history reconstruction for the 20th century. *Global Change Biology*, **11**, 398–420.
- Ohlrogge J, Allen D, Berguson B, DellaPenna D, Shachar-Hill Y, Szyme S (2009) Driving on biomass. *Science*, **324**, 1019–1020.
- Owens AK, Moseley KR, McCay TS, Castleberry SB, Kilgo JC, Ford WM (2008) Amphibian and reptile community response to coarsewoody debris manipulations in upland loblolly pine (*Pinus taeda*) forests. *Forest Ecology and Management*, **256**, 2078–2083.
- Pausas JG, Keely JE (2009) A burning story: the role of fire in the history of life. *BioScience*, **59**, 593–601.
- Pechony O, Shindell DT (2010) Driving forces of global wildfires over the past millennium and the forthcoming century. *Proceedings of the National Academy of Sciences*, **107**, 19167–19170.
- Piñeiro G, Jobbagy EG, Baker J, Murray BC, Jackson RB (2009) Set-asides can be better climate investment than corn ethanol. *Ecological Applications*, **19**, 277–282.
- Potter CS, Randerson JT, Field CB, Matson PA, Vitousek PM, Mooney HA, Klooster SA (1993) Terrestrial ecosystem production: a process model based on global satellite and surface data. *Global Biogeochemical Cycles*, **7**, 811–841.
- Preisler HK, Westerling AL, Gebert KM, Munoz-Arriola F, Holmes TP (2011) Spatially explicit forecasts of large wildland fire probability and suppression costs for California. *International Journal of Wildland Fire*, **20**, 508–517.
- Ramanathan V, Carmichael G (2008) Global and regional climate changes due to black carbon. *Nature Geoscience*, **1**, 221–227.
- Richter DD Jr, Jenkins DH, Karakash JT, Knight J, McCreery LR, Nemesthoy KP (2009) Wood energy in America. *Science*, **323**, 1432–1433.
- Saugier B, Roy J, Mooney HA (2001) Estimations of global terrestrial productivity: converging toward a single number? In: *Terrestrial Global Productivity* (eds Roy J, Saugier B, Mooney HA), pp. 543–556. Academic Press, San Diego.
- Schiermeier Q, Tollefson J, Scully T, Witze A, Morton O (2008) Electricity without carbon. *Nature*, **454**, 816–823.
- Schroeder W, Csizsar I, Giglio L, Schmidt CC (2010) On the use of fire Radiative power, area, and temperature estimates to characterize biomass burning via moderate to coarse spatial resolution remote sensing data in the Brazilian Amazon. *Journal of Geophysical Research*, **115**, D21121, doi: 10.1029/2009JD013769.
- Searchinger T, Heimlich R, Houghton RA *et al.* (2008) Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land use change. *Science*, **319**, 1238–1240.
- Skinner WR, Shabbar A, Flannigan MD, Logan K (2006) Large forest fires in Canada and the relationship to global sea surface temperatures. *Journal of Geophysical Research*, **111**, D14106.
- Smil V (2003) *Energy at the Crossroads: Global Perspectives and Uncertainties*. The MIT Press, Cambridge, MA.
- Tymstra C, Flannigan MD, Armitage OB, Logan KA (2007) Impact of climate change on area burned in Alberta's boreal forest. *International Journal of Wildland Fire*, **16**, 153–160.
- Vázquez DP, Alvarez JA, Debandi G, Aranibar JN, Villagra PE (2012) Ecological consequences of dead wood extraction in an arid ecosystem. *Basic and Applied Ecology*, **12**, 722–732.
- van der Werf GR, Dempewolf J, Trigg SN *et al.* (2008) Climate regulation of fire emissions and deforestation in equatorial Asia. *Proceedings of the National Academy of Sciences*, **105**, 20350–20355.
- van der Werf GR, Randerson JT, Giglio L *et al.* (2010) Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmospheric Chemistry and Physics*, **10**, 11707–11735.
- Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW (2006) Warming and earlier spring increase Western U.S. forest wildfire activity. *Science*, **313**, 940–943.
- Westerling AL, Turner MG, Smithwick EAH, Romme WH, Ryan MG (2011) Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. *Proceedings of the National Academy of Sciences*, **108**, 13165–13170.
- Wooster MJ, Roberts G, Perry GLW, Kaufman YJ (2005) Retrieval of biomass combustion rates and totals from fire radiative power observations: FRP derivation and calibration relationships between biomass consumption and fire radiative energy release. *Journal of Geophysical Research*, **110**, D24311.
- Wotton BM, Nock CA, Flannigan MD (2010) Forest fire occurrence and climate change in Canada. *International Journal of Wildland Fire*, **19**, 253–271.
- Yagi K, Nakata T (2011) Economic analysis on small-scale forest biomass gasification considering geographical resources distribution and technical characteristics. *Biomass and Bioenergy*, **35**, 2883–2892.
- Zobler L (1986) *A world soil file for global climate modeling*. NASA Tech. Memo. 87802. NASA/GISS, New York.

**Supporting Information**

Additional Supporting Information may be found in the online version of this article:

**Table S1.** Ecological and economic variables used to explore bioelectricity opportunities at across countries.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.