

# CO<sub>2</sub> enrichment and soil type additively regulate grassland productivity

H. Wayne Polley<sup>1</sup> , Michael J. Aspinwall<sup>2,3</sup> , Harold P. Collins<sup>1</sup>, Anne E. Gibson<sup>1</sup>, Richard A. Gill<sup>4</sup> , Robert B. Jackson<sup>5</sup> , Virginia L. Jin<sup>6</sup> , Albina R. Khasanova<sup>2</sup> , Lara G. Reichmann<sup>2,7</sup>  and Philip A. Fay<sup>1</sup> 

<sup>1</sup>Agricultural Research Service, Grassland, Soil and Water Research Laboratory, USDA, 808 East Blackland Road, Temple, TX 76502, USA; <sup>2</sup>Section of Integrative Biology, The University of Texas at Austin, 1 University Station C0930, Austin, TX 78712, USA; <sup>3</sup>Department of Biology, University of North Florida, 1 UNF Drive, Jacksonville, FL 32224, USA; <sup>4</sup>Department of Biology, Brigham Young University, 4102 LSB, Provo, UT 84602, USA; <sup>5</sup>Department of Earth System Science, Woods Institute for the Environment and Precourt Institute for Energy, Stanford University, Y2E2 Building, 379B, Stanford, CA 94305, USA; <sup>6</sup>Agricultural Research Service, ARS Agroecosystem Management Research Unit, USDA, University of Nebraska, 251 Filley Hall, Lincoln, NE 68583, USA; <sup>7</sup>Data Institute, University of San Francisco, 101 Howard St., San Francisco, CA 94105, USA

## Summary

Author for correspondence:

H. Wayne Polley

Tel: +1 254 770 6629

Email: wayne.polley@ars.usda.gov

Received: 2 May 2018

Accepted: 18 October 2018

*New Phytologist* (2019) **222**: 183–192

doi: 10.1111/nph.15562

**Key words:** aboveground productivity, atmospheric CO<sub>2</sub> enrichment, rangeland, soil texture, soil water potential, species composition, subambient CO<sub>2</sub> concentration.

- Atmospheric CO<sub>2</sub> enrichment usually increases the aboveground net primary productivity (ANPP) of grassland vegetation, but the magnitude of the ANPP–CO<sub>2</sub> response differs among ecosystems. Soil properties affect ANPP via multiple mechanisms and vary over topographic to geographic gradients, but have received little attention as potential modifiers of the ANPP–CO<sub>2</sub> response.
- We assessed the effects of three soil types, sandy loam, silty clay and clay, on the ANPP response of perennial C<sub>3</sub>/C<sub>4</sub> grassland communities to a subambient to elevated CO<sub>2</sub> gradient over 10 yr in Texas, USA. We predicted an interactive, rather than additive, effect of CO<sub>2</sub> and soil type on ANPP.
- Contrary to prediction, CO<sub>2</sub> and soil additively influenced grassland ANPP. Increasing CO<sub>2</sub> by 250 μl l<sup>-1</sup> increased ANPP by 170 g m<sup>-2</sup> across soil types. Increased clay content from 10% to 50% among soils reduced ANPP by 50 g m<sup>-2</sup>. CO<sub>2</sub> enrichment increased ANPP via a predominant direct effect, accompanied by a smaller indirect effect mediated by a successional shift to increased dominance of the C<sub>4</sub> tallgrass *Sorghastrum nutans*.
- Our results indicate a large, positive influence of CO<sub>2</sub> enrichment on grassland productivity that resulted from the direct physiological benefits of CO<sub>2</sub> augmented by species succession, and was expressed similarly across soils of differing physical properties.

## Introduction

The development of a predictive understanding of how atmospheric CO<sub>2</sub> enrichment is affecting the primary productivity of the terrestrial biosphere is among the most pressing of ecological challenges. The terrestrial biosphere absorbs *c.* 25% of anthropogenic carbon (C) emissions (Le Quéré *et al.*, 2018). Uncertainty in CO<sub>2</sub> effects on ecosystem C uptake is a major constraint in the prediction of C cycling and the provisioning of productivity-related ecosystem services.

Grasslands cover *c.* 25% of the terrestrial area and are an important contributor to the global C balance (Sala *et al.*, 1996). CO<sub>2</sub> enrichment stimulates the aboveground net primary productivity (ANPP) of most water-limited grasslands by increasing plant water use efficiency (WUE; productivity per unit of transpiration; Morgan *et al.*, 2004; Nowak *et al.*, 2004; Fatichi *et al.*, 2016), but grassland ANPP, as other ecosystem functions, is determined by drivers in addition to water availability which act simultaneously and often interactively with CO<sub>2</sub> (Polley *et al.*,

2011). CO<sub>2</sub> enrichment usually shows greater stimulation of plant productivity when nitrogen (N) availability is relatively high (Owensby *et al.*, 1994; Reich & Hobbie, 2013; Mueller *et al.*, 2016), for example. Other drivers include precipitation timing (Hovenden *et al.*, 2014), disturbance regimes (Newton *et al.*, 2014), plant species composition (Langley & Megonigal, 2010; Fay *et al.*, 2012; Polley *et al.*, 2012) and soil properties (Epstein *et al.*, 1997, 1998), including soil texture, which influences water availability to plants (Tor-Ngern *et al.*, 2017).

Soils have received little attention as potential mediators of the ANPP–CO<sub>2</sub> response, apart from the role of soil properties in influencing N and water availability to plants. Spatial variation in soil N is common (García-Palacios *et al.*, 2012), with possible feedbacks on the composition, productivity and CO<sub>2</sub> response of plant assemblages (Owensby *et al.*, 1994; Niklaus & Körner, 2004; Langley & Megonigal, 2010; Reich & Hobbie, 2013; Mueller *et al.*, 2016). Water availability varies across gradients in soil physical and related hydrological properties at spatial scales over which precipitation is similar. CO<sub>2</sub> enrichment typically

shows greater enhancement of productivity in drier years or at drier localities, the result of increased WUE (Owensby *et al.*, 1999; Morgan *et al.*, 2004; Nowak *et al.*, 2004; Fatichi *et al.*, 2016). A similar mechanism may contribute to soil differences in the productivity response to CO<sub>2</sub> enrichment. By increasing WUE, CO<sub>2</sub> enrichment could lead to a greater increase in plant productivity on fine- than coarse-textured soils, because soil water availability to plants (soil water potential) at a given level of extractable soil water is usually lower in fine- than coarse-textured soils (Tor-Ngern *et al.*, 2017).

Soils can also mediate the ANPP–CO<sub>2</sub> response by altering the plant species composition or abundance (Polley *et al.*, 2011; Fay *et al.*, 2015a). Soils with a high percentage of sand content favor productive C<sub>4</sub> tall grasses (Epstein *et al.*, 1998), which may experience greater benefits from CO<sub>2</sub> enrichment relative to shorter statured grasses, for example (Fay *et al.*, 2012; Polley *et al.*, 2012).

The response of ANPP to CO<sub>2</sub> enrichment is likely to differ among soil types when soil effects on productivity are tightly coupled with soil differences in resource (e.g. water, N) availability. Soil-specific responses to CO<sub>2</sub> enrichment are indicated statistically by a soil type × CO<sub>2</sub> interaction. However, soil physical properties may affect the ANPP–CO<sub>2</sub> response via mechanisms that are independent of, or weakly correlated with, soil differences in resource availability. Plant growth and species abundances are sensitive to variation in soil strength or compaction (Masle & Passioura, 1987; Wallace, 1987; Schrama *et al.*, 2013), for example. In these cases, soil and CO<sub>2</sub> effects on productivity may be additive, the sum of their separate effects, rather than interactive (Zavaleta *et al.*, 2003). Additive effects are evidenced by the absence of a statistical CO<sub>2</sub> × soil interaction.

We assessed the effects of soil type and CO<sub>2</sub> on the ANPP of perennial grassland communities during a 10-yr field experiment on three soil types in Texas, USA (Polley *et al.*, 2008; Fay *et al.*, 2009). Identical assemblages of C<sub>3</sub> forbs and C<sub>4</sub> grasses were established on three soil types (clay, sandy loam, silty clay), representative of the variation in soil texture and associated hydrological properties in the Blackland Prairie region of Texas. Communities on each soil type were grown along a continuous CO<sub>2</sub> gradient from the pre-industrial concentration (250 μl l<sup>-1</sup>) to a level expected this century (500 μl l<sup>-1</sup>).

The ANPP–CO<sub>2</sub> response differed among soils during the initial 5 yr of this experiment (Fay *et al.*, 2012; Polley *et al.*, 2012). CO<sub>2</sub> enrichment directly increased ANPP on all soil types during at least one of the first 5 yr, probably by increasing leaf photosynthesis rates and leaf-level photosynthetic WUE (Fay *et al.*, 2012). CO<sub>2</sub> effects differed among soils during these initial years because CO<sub>2</sub> feedbacks on productivity through changes in the soil water potential (Fay *et al.*, 2012) and relative abundance of a productive C<sub>4</sub> grass, *Sorghastrum nutans* (Fay *et al.*, 2012; Polley *et al.*, 2012), differed in magnitude among soils. It remains unresolved as to whether the processes that contributed initially to interactive soil × CO<sub>2</sub> effects on productivity are sustained over longer time periods. N availability did not differ among soils initially (Fay *et al.*, 2012), but longer term changes in N availability (Reich *et al.*, 2018) could alter the relative roles of species abundances,

soil water and N in mediating soil effects on the ANPP–CO<sub>2</sub> response.

We predicted that [CO<sub>2</sub>] effects on ANPP would continue to differ among soil types over 10 yr of CO<sub>2</sub> enrichment. Specifically, we predicted first that CO<sub>2</sub> enrichment would elicit the largest absolute increase in ANPP on soils for which mean ANPP was greatest, consistent with evidence that CO<sub>2</sub> leads to a greater stimulation of plant productivity when N availability is high rather than low (Owensby *et al.*, 1994; Reich & Hobbie, 2013; Mueller *et al.*, 2016). Second, we predicted that CO<sub>2</sub> enrichment would elicit the largest proportional increase in ANPP on the soil with the lowest mean water potential, consistent with evidence that CO<sub>2</sub> leads to a greater stimulation of plant productivity when plant growth is water limited (Morgan *et al.*, 2004; Nowak *et al.*, 2004; Fatichi *et al.*, 2016). Third, we predicted that initial, soil-specific responses to CO<sub>2</sub> would be amplified over time as the result of a progressive increase in soil differences in *Sorghastrum* dominance at elevated CO<sub>2</sub>. Progressive CO<sub>2</sub> effects on soil resource levels (Luo *et al.*, 2004; Reich & Hobbie, 2013) or plant composition or functioning (Niklaus *et al.*, 2001; Hungate *et al.*, 2004) may reinforce interactive controls on the productivity–CO<sub>2</sub> response. However, the initial soil × CO<sub>2</sub> interaction for ANPP could be weakened or even eliminated if changes in N mineralization rates (Reich *et al.*, 2018) or climatic variation (Zhu *et al.*, 2016) reverse progressive CO<sub>2</sub> effects that reinforce soil-specific ANPP–CO<sub>2</sub> responses. Answers to questions as to whether or how soil type influences the CO<sub>2</sub> response of productivity will improve our understanding and modeling of grassland responses to the historical and anticipated rise in CO<sub>2</sub> concentration across edaphically variable landscapes.

## Materials and Methods

### Facility

We used the Lysimeter CO<sub>2</sub> Gradient (LYCOG) facility located outdoors in central Texas, USA to expose grassland communities to a continuous gradient in CO<sub>2</sub> spanning pre-industrial to elevated concentrations (250–500 μl l<sup>-1</sup>) each growing season for a decade (2006–2015) (Supporting Information Fig. S1; Johnson *et al.*, 2000; Polley *et al.*, 2008; Fay *et al.*, 2009, 2015b). LYCOG consists of two transparent, tunnel-shaped chambers, aligned parallel along a north–south axis. Each chamber is divided into 10 consecutive compartments, each 5 m long and 1.2 m wide. The aerial portion of each compartment was built atop a 5-m-long steel container that had been buried to a depth of 1.2 m. The aerial growth of the chambered vegetation was enclosed in a transparent polyethylene film. The desired [CO<sub>2</sub>] gradients were maintained by automatically varying the rate of air flow through the chambers in response to changes in photosynthesis (daylight) or respiration (night) rates. Night-time [CO<sub>2</sub>] was regulated at 130–150 μl l<sup>-1</sup> above daytime values along each chamber. Air temperature in the chambers was regulated with the goal of matching ambient values by cooling (and dehumidifying) air at 5-m intervals along the chambers. This goal was achieved most consistently during the cooler spring and

autumn months. The chamber air temperature was  $1.87 \pm 0.33^\circ\text{C}$  cooler than the ambient temperature on average across the full growing season. The air temperature, dew point and  $[\text{CO}_2]$  were measured at the air entrance and exit of each 5-m compartment at 20-min intervals.  $\text{CO}_2$  treatments were maintained each growing season (2006–2015) from March/April through mid-November.

### Plant communities

Communities of grassland species were grown in intact soil monoliths (each 1 m  $\times$  1 m  $\times$  1.5 m deep). Monoliths of three soil types were collected in the Texas Blackland Prairie region (Austin silty clay,  $n=32$ ; Houston Black clay,  $n=32$ ; Bastisil sandy loam,  $n=16$ ). Monoliths were collected by pressing open-ended steel boxes into the soil. Steel-encased monoliths were placed in containers beneath 5-m-long compartments. Each compartment contained two monoliths of each of two of the three soil types. The monolith order within compartments was assigned randomly. Monoliths of the sandy loam soil were included in the second of each consecutive pair of compartments along the chambers. Seedlings of seven perennial species, including four  $\text{C}_4$  grasses, two  $\text{C}_3$  forbs and one herbaceous legume, were transplanted into each monolith 3 yr before  $\text{CO}_2$  treatment. All species are native to central Texas (Diggs *et al.*, 1999) and present in the Texas Blackland Prairie. Eventual community dominants included the  $\text{C}_4$  grasses *Bouteloua curtipendula* (Michx.) Torr. and *Sorghastrum nutans* (L.) Nash, and the forb species *Solidago canadensis* L. No fertilizer was applied. Grubs (Coleoptera: Scarabaeidae) killed most transplants in 20 of the total of 80 monoliths (eight silty clay, 12 clay) during the first year of  $\text{CO}_2$  treatment. These monoliths were replanted in spring 2007 as *Panicum virgatum* L. (switchgrass) monocultures that were well watered and fertilized for the duration of the experiment to facilitate  $\text{CO}_2$  regulation by increasing the photosynthetic sink. Sixty monoliths of species mixtures remained (silty clay,  $n=24$ ; clay,  $n=20$ ; sandy loam,  $n=16$ ). We excluded data from one silty clay monolith on which most vegetation died following year 3 of  $\text{CO}_2$  regulation.

### Implementation

We surface irrigated each monolith with the equivalent of the growing season (April–October) average of precipitation at our site (560 mm) during each season. We varied the fraction of growing season irrigation that was added in spring vs summer among years to create interannual differences in seasonal water availability, whilst holding the annual irrigation amount constant. All monoliths were subjected to the same irrigation schedule each year. The spring (mid-April through mid-June) fraction of total irrigation that was applied to each of the 60 monoliths with species mixtures was varied interannually between 19 and 43%, the latter being the long-term mean for the site (range = 10–80%; 100-yr record; Fig. S2). Interannual differences in the seasonality of irrigation combined with soil

hydrological differences and seasonal or interannual variation in plant transpiration to create variation in plant-available soil water.

Daytime  $[\text{CO}_2]$  varied as a linear function of distance along the chambers (Fay *et al.*, 2009). The mean  $[\text{CO}_2]$  to which each monolith was exposed was calculated using linear regressions of  $[\text{CO}_2]$  vs distance. Calculated  $[\text{CO}_2]$  was used as the independent variable in regression analyses with ANPP and species abundances.

### Measurements

Volumetric soil water content (vSWC) was measured weekly in the center of each monolith with a neutron probe. Soil water (matric) potential over a depth of 0–50 cm ( $\Psi_s$ ) was calculated from vSWC using soil-specific  $\Psi_s$ –vSWC relationships developed from measurements of  $\Psi_s$  with a dewpoint hygrometer (Model WP4T; Decagon Devices, Pullman, WA, USA) (Polley *et al.*, 2016).

Soil N availability was assessed in 2008, 2010, 2013 and 2015 (years 3, 5, 8 and 10 of  $\text{CO}_2$  treatment) using ion exchange resins (PRS-probes; Western Ag Innovations, Saskatoon, SK, Canada) to measure  $\text{NO}_3^-$  and  $\text{NH}_4^+$  in soil solution. Resin probes were installed monthly during each of 2 months in both spring (mid-April to mid-June) and summer (mid-July to mid-September) each year. N amounts were measured colorimetrically for probes combined from the two monoliths of a given soil type in each 5-m-long compartment of the  $\text{CO}_2$  chambers. Soil N availability per sampling location and year was taken as the monthly average of  $\mu\text{g N}$  accumulated per  $10\text{ cm}^2$  of probe surface area.

All vegetation from each monolith was clipped at a height of 5 cm at the end of each growing season following plant senescence. Clipped vegetation was dried ( $65^\circ\text{C}$ ) and weighed by species. Species abundances were calculated as a fraction of total ANPP. Mean ANPP of monoliths was similar in magnitude to the productivity measured in two remnant prairies (clay/silty clay soils) over 5 yr (mean = 309 and  $379\text{ g m}^{-2}\text{ yr}^{-1}$ ; Polley *et al.*, 2007), implying no discernible limitation of monolith rooting volume on plant growth.

### Data analysis

We used a repeated-measures analysis of covariance (PROC MIXED in SAS 9.4) to model treatment effects on ANPP, fractional contribution of the tall grass *Sorghastrum* to community productivity (fSorg),  $\Psi_s$  and soil N:

$$y = \text{intercept} + \text{CO}_2 + \text{soil} + \text{monolith}(\text{soil}) \\ + \text{compartment}(\text{soil}) + \text{chamber}(\text{compartment}) + (\text{CO}_2 \\ \times \text{soil}) + \text{year} + (\text{CO}_2 \times \text{year}) + (\text{soil} \times \text{year}) + e$$

Eqn 1

where  $\text{CO}_2$  was the covariate, soil was a fixed effect, monoliths within soil type (monolith(soil)), soil types within 5-m-long compartments (compartment(soil)) and compartments within

chambers (chambers(compartment)) were random effects,  $\text{CO}_2 \times \text{soil}$  was the soil-specific covariate and  $e$  was the random error. Year was a repeated effect fitted using an autoregressive covariance structure.  $\text{CO}_2 \times \text{year}$  and  $\text{soil} \times \text{year}$  terms modeled interannual variation in  $\text{CO}_2$  and soil effects on dependent variables.

We also fitted Structural Equation Models (SEMs) to evaluate causal relationships among interacting variables that together affected ANPP (Shipley, 2000; Grace, 2006). Models were fitted using IBM SPSS AMOS 21 software. Our initial (meta) model of predicted ANPP ( $\log_e(\text{ANPP})$ ) included two independent variables,  $[\text{CO}_2]$  and soil type, the latter represented as the percentage clay content ( $\log_e(\% \text{ clay})$ ), and three dependent variables,  $\Psi_s$ ,  $f\text{Sorg}$  and a lagged mediator of  $f\text{Sorg}$ , the *Sorghastrum* fraction of productivity during the previous year ( $f\text{Sorg}_p$ ). Soil N was not included in the SEM as N did not vary with  $[\text{CO}_2]$  or soil type and was weakly correlated with ANPP (see Results). SEMs were fitted to a dataset that included measurements from all monoliths, soil types and 10 yr, and separately to data from the first 5 yr or final 5 yr of the experiment. We iteratively removed parameters or pathways from the meta model in order to identify models that yielded the most parsimonious fit among those tested, as indicated by smaller values of the root-mean-square error of approximation (RMSEA) and Akaike's Information Criterion (AIC).

#### Data accessibility


Data have been deposited in the Dryad Digital Repository doi:10.5061/dryad.1hc582s.

## Results

### Soil and $\text{CO}_2$ effects on biotic and abiotic variables


Grasses dominated the vegetation on all soils, but species abundances differed among soil types (Fig. 1).  $\text{C}_4$  grasses contributed an average of 73%, 84% and 88% of annual productivity on clay, silty clay and sandy loam soils, respectively, over 10 yr. *Sorghastrum nutans* was the dominant grass on the clay and sandy loam soils (32% and 46% of ANPP, respectively) and subdominant to *Bouteloua curtipendula* on the silty clay soil, where *Sorghastrum* contributed 25% of ANPP. *Solidago canadensis* was the dominant  $\text{C}_3$  forb, contributing a mean of 9%, 10% and 15% of annual ANPP on the sandy loam, silty clay and clay soils, respectively, over 10 yr. Because grasses dominated communities, ANPP was largely dependent on the growth of these  $\text{C}_4$  species.

$\text{CO}_2$  enrichment increased aboveground productivity across soil types (Table S1; Fig. 2). The  $[\text{CO}_2]$  effect on ANPP did not differ significantly among soils (analysis of covariance;  $\text{CO}_2 \times \text{soil}$ ,  $P=0.12$ ) (Table S1), but varied among years. The slopes of ANPP– $\text{CO}_2$  relationships were significantly smaller in 2011 and 2013 (years 6 and 8) than in the remaining years (50.8 vs 114.0  $\text{g m}^{-2}$  increase in ANPP per 100  $\mu\text{l l}^{-1}$  increase in  $[\text{CO}_2]$ ; not shown).



**Fig. 1** Mean ( $\pm$  SE) values of aboveground net primary productivity (ANPP) for grassland communities (white circles) and component  $\text{C}_4$  grasses (gray squares) and  $\text{C}_3$  forb species (closed circles) for each of three soil types and year of  $\text{CO}_2$  treatment. Values are averaged across  $[\text{CO}_2]$  treatments for each year and soil ( $n=16\text{--}23$ ). \*in (b) indicates years in which ANPP was not significantly lower for the silty clay than clay and sandy loam soils.

ANPP was greater for communities grown on clay and sandy loam soils than on silty clay soil over 10 yr (Fig. 1; mean = 478, 462 and 331  $\text{g m}^{-2} \text{ yr}^{-1}$ , respectively), although soil differences in ANPP varied among years (Table S1). ANPP was smaller on the silty clay than on both the clay and sandy loam soils in all but



**Fig. 2** Mean ( $\pm$  SE) responses for total aboveground net primary productivity (ANPP) of communities and the contribution of the tall grass *Sorghastrum* to community productivity (fraction *Sorghastrum*) along a [CO<sub>2</sub>] gradient for each of three soil types over a decade of CO<sub>2</sub> treatment. Lines show fits of linear or quadratic regressions to 10-yr averages per monolith for sandy loam (shaded squares; adj.  $r^2 = 0.53$  (a) and 0.19 (d);  $P < 0.05$ ), silty clay (open circles; adj.  $r^2 = 0.53$  (b) and 0.39 (e);  $P < 0.0007$ ) and clay soils (closed circles, adj.  $r^2 = 0.36$ ,  $P = 0.02$  (c) and not significant (f)).


two (2007, 2011) of the first 8 yr, and on both the silty clay and sandy loam soils than on the clay soil during the final year (Fig. 1).

CO<sub>2</sub> enrichment increased ANPP partly by favoring the dominant grass species. Increased [CO<sub>2</sub>] boosted the *Sorghastrum* fraction of ANPP (fSorg) (Table S1; Fig. 2). Increased relative abundance of *Sorghastrum*, in turn, increased ANPP across years for each soil type (Fig. 3). fSorg explained 24% of the variance in ANPP across soils and years (ANPP (g m<sup>-2</sup>) = 332.2 + 274.1 × fSorg;  $P < 0.0001$ ). The CO<sub>2</sub> effect on fSorg changed with time. The contribution of *Sorghastrum* to community ANPP was greater when averaged across elevated [CO<sub>2</sub>] than subambient [CO<sub>2</sub>] beginning in 2007, year 2 ((fSorg (E-S) > 0); Fig. 4), especially on silty clay and sandy loam soils. The strong initial response of *Sorghastrum* to CO<sub>2</sub> was transient. fSorg(E-S) declined on all three soils beginning in 2010–2011 (years 5–6), with the largest declines in 2012–2013 (years 7–8). The *Sorghastrum* fraction of ANPP (fSorg) did not differ significantly among soils (Table S1); however, nor did soil influence the [CO<sub>2</sub>] effect on fSorg.


[CO<sub>2</sub>] and soil effects on soil water potential exerted minor feedback on ANPP. CO<sub>2</sub> enrichment increased  $\Psi_s$  to 50 cm depth, most strongly for the clay soil, by 0.4 MPa per 100 μl l<sup>-1</sup> rise in [CO<sub>2</sub>] (Fig. 5; CO<sub>2</sub> × soil,  $P = 0.02$ ; Table S1).  $\Psi_s$  differed among soils ( $P < 0.0001$ ). Soil water availability was highest for the sandy loam, intermediate for the silty clay and lowest for

the clay (−0.27, −0.73 and −1.76 MPa, respectively).  $\Psi_s$  differed among soils because soils differed in hydrological properties that influenced relationships between  $\Psi_s$  and vSWC. The range in vSWC created by irrigation treatments and plant transpiration traversed a portion of the  $\Psi_s$  vs vSWC curve over which the variation in  $\Psi_s$  was relatively large in the clay soil. Conversely, annual means of vSWC did not decline to levels at which  $\Psi_s$  responded greatly to the change in water content in the sandy loam soil. Soil differences in  $\Psi_s$  vs vSWC relationships are consistent with differences expected from soil variation in texture (% clay content = 9.4%, 43.1% and 52.6% for the sandy loam, silty clay and clay soils, respectively). Over the range of  $\Psi_s$  encountered, water potential alone explained little (< 1%) of the variance in ANPP across data from soils and years combined (ANPP = 397.0 − 17.0 ×  $\Psi_s$ ;  $P = 0.08$ ; Fig. 3).

Resin-available soil N, by contrast, did not vary with [CO<sub>2</sub>] or differ among soil types over 4 yr of N measurements (Table S1). Soil N declined from 8.3 ± 0.3 μg N 10 cm<sup>-2</sup> of probe area month<sup>-1</sup> in 2008 (year 3) to 3.0 ± 0.3 μg N 10 cm<sup>-2</sup> month<sup>-1</sup> in 2010 (year 5), when ANPP was near maximal on all soils (Fig. 1), and remained low thereafter (2.3 ± 0.1 and 1.4 ± 0.1 μg N 10 cm<sup>-2</sup> month<sup>-1</sup> in years 8 and 10, respectively). Neither fSorg nor  $\Psi_s$  was correlated with soil N ( $P = 0.94$  and 0.11, respectively). ANPP was a weak positive function of resin-available N across years, soils and [CO<sub>2</sub>] (not shown; adj.  $r^2 = 0.03$ ;  $P = 0.03$ ), but was not significantly correlated with the



**Fig. 3** Relationships between aboveground net primary productivity (ANPP) of grassland communities and (a–c) the contribution of the tall grass *Sorghastrum* to community productivity (fraction *Sorghastrum*) and (d–f) the growing season mean of the soil water potential to 50 cm depth for each of three soil types over a decade of CO<sub>2</sub> treatment. Lines show fits of linear (a–c, e) or exponential (d) regressions to the data from all CO<sub>2</sub> treatments, monoliths and years for sandy loam (shaded squares; adj.  $r^2 = 0.15$  and  $0.21$ ,  $P < 0.0001$ ,  $n = 160$  (a, d)), silty clay (open circles; adj.  $r^2 = 0.24$  and  $0.01$ ,  $P < 0.0001$  and  $0.06$ ,  $n = 230$  (b, e)) and clay soils (closed circles, adj.  $r^2 = 0.20$ ,  $P < 0.0001$  and  $P = 0.52$ ,  $n = 200$  (c, f)). The scale of the x-axis differs among panels (d)–(f).




**Fig. 4** Temporal trends in the CO<sub>2</sub> effect on the relative abundance of the tallgrass species *Sorghastrum* for grassland communities grown on each of three soil types. Shown are the annual values of the elevated vs subambient difference (E–S) in fractional contribution of *Sorghastrum* to community aboveground net primary productivity (ANPP) for silty clay (SC; open circles), sandy loam (SL; gray squares) and clay soils (C; closed circles). The inset shows 10-yr averages of the E–S difference in *Sorghastrum* fraction for each soil type.

locations of 5-m-long compartments along chambers (chamber position) after accounting for effects of variation in [CO<sub>2</sub>] and fSorg using a sequential regression analysis ( $P > 0.15$ ).

#### Structural equation models

SEMs were used to test joint effects of the two independent variables, [CO<sub>2</sub>] and log<sub>e</sub>(% clay), and three dependent variables,  $\Psi_s$ , fSorg and fSorg<sub>p</sub>, on ANPP (log<sub>e</sub>(ANPP)). This model adequately fitted the data from all soils, monoliths and years combined and explained 37% of the variance in ANPP (Fig. 6). Both the [CO<sub>2</sub>] and log<sub>e</sub>(% clay) effects on ANPP were primarily direct, rather than mediated through changes in fSorg or  $\Psi_s$ . The direct CO<sub>2</sub> effect was positive and accounted for 80% of the total CO<sub>2</sub> effect on ANPP. The direct effect of soil type (clay content) on ANPP was negative. CO<sub>2</sub> enrichment indirectly increased ANPP by increasing fSorg, but reduced ANPP by increasing  $\Psi_s$ , because mean ANPP was greater for the clay soil with lowest mean  $\Psi_s$  than the silty clay soil with higher potential. Indirect effects of greater clay content on ANPP were opposite in sign to those of [CO<sub>2</sub>]. The total positive effect of CO<sub>2</sub> enrichment was twice the negative effect of clay content. Increasing CO<sub>2</sub> by 100  $\mu\text{l l}^{-1}$  increased ANPP by 85 g m<sup>-2</sup>, whereas increasing the clay content by 10% reduced ANPP by 12 g m<sup>-2</sup>.




**Fig. 5** Relationships between the soil water potential to 50 cm depth ( $\Psi_s$ ) of three soil types and three variables: atmospheric  $[\text{CO}_2]$  (a), soil percentage clay content (b) and volumetric water content (vSWC; c). Lines in (a) show linear regression fits to 10-yr averages per monolith ( $\pm$  SE) for sandy loam (adj.  $r^2 = 0.26$ ,  $P = 0.04$ ,  $n = 14$  after deleting two outlier points), silty clay (adj.  $r^2 = 0.10$ ,  $P = 0.08$ ,  $n = 23$ ) and clay (adj.  $r^2 = 0.31$ ,  $P = 0.006$ ,  $n = 20$ ) soils. Shown in (b) are the growing season means of  $\Psi_s$  for each of the 10 yr of  $\text{CO}_2$  treatment. Brackets in (c) denote the range in growing season means of vSWC observed for each soil type.  $\Psi_s$  was calculated from measurements of vSWC using soil-specific  $\Psi_s$ -vSWC relationships developed using a dewpoint hygrometer.

The predominant pathways via which  $[\text{CO}_2]$  and soil type influenced ANPP differed between SEMs fitted to the data from the first 5 yr vs final 5 yr of the experiment (Figs S3, S4). The soil effect on ANPP was entirely indirect for the first 5 yr of the experiment (2006–2010; Fig. S3), mediated through fSorg. By contrast, neither  $[\text{CO}_2]$  nor the percentage clay content influenced the current year *Sorghastrum* fraction (fSorg) via a direct pathway during the final 5 yr of the experiment (2011–2015; Fig. S4).

## Discussion

Contrary to our prediction that the ANPP- $\text{CO}_2$  response would differ among soils, we found that atmospheric  $[\text{CO}_2]$  and soil type additively affected the aboveground productivity of perennial grassland communities. Increasing  $\text{CO}_2$  from 280 to 480  $\mu\text{l l}^{-1}$  increased productivity by *c.* 170  $\text{g m}^{-2}$  across soil types, a productivity enhancement comparable with that anticipated from a 290-mm increase in mean annual precipitation (MAP) across grasslands in the North American Central Plains (Sala *et al.*, 1988). Conversely, increasing clay content from 10% (sandy loam soil) to 50% (clay soil) reduced productivity by *c.* 50  $\text{g m}^{-2}$ . Also contrary to prediction,  $\text{CO}_2$  enrichment did not elicit a larger absolute increase in ANPP on soils with greatest mean ANPP (sandy loam, clay) or a larger relative increase in ANPP on the soil with the lowest mean water potential (clay). Initial soil differences in the ANPP- $\text{CO}_2$  response were also not amplified in later years by continued soil-specific increases in *Sorghastrum* dominance at elevated  $\text{CO}_2$ . Our results indicate a large positive influence of  $\text{CO}_2$  enrichment from subambient to elevated concentrations on grassland productivity which, under the conditions of this experiment, was expressed similarly across soil types that differed in physical and related hydrological properties.

Three factors may explain why initial soil-specific effects on the ANPP- $\text{CO}_2$  response were not maintained over 10 yr. First,  $\text{CO}_2$  effects on ANPP were primarily direct rather than mediated by soil differences in  $\text{CO}_2$  effects on soil water or N.  $\text{CO}_2$  and soil effects on water and N availability were apparently too small to significantly influence the ANPP- $\text{CO}_2$  response. Second,  $\Psi_s$  did not consistently regulate productivity across soil types. Water limitation frequently enhances the relative effect of  $\text{CO}_2$  enrichment on productivity (Morgan *et al.*, 2004; Nowak *et al.*, 2004; Fatichi *et al.*, 2016), although this result is not universal (Dukes *et al.*, 2005; Hovenden *et al.*, 2014). In this experiment, however,  $\Psi_s$  apparently did not decline to levels sufficient to consistently constrain growth across soil types. Third, the progressive increase in *Sorghastrum* abundance at higher  $\text{CO}_2$ , which amplified soil differences in the ANPP- $\text{CO}_2$  response during the initial 3–5 yr of treatment (Fay *et al.*, 2012), ended following year 6, resulting in only a small *Sorghastrum*-mediated effect of  $\text{CO}_2$  enrichment on ANPP over 10 yr. Year 6 (2011) was characterized by the most severe drought in Texas in a record extending to 1895 and was accompanied by elevated temperatures and evaporative demand (Hoerling *et al.*, 2013). Growing season temperatures in  $\text{CO}_2$  chambers were higher in 2011 than during the 4 yr that



**Fig. 6** Structural equation model of variables affecting the annual aboveground net primary productivity [ $\log_e(ANPP)$ ] of grassland communities over a decade ( $\chi^2 = 0.61$ ,  $df = 3$ ,  $P = 0.90$ ,  $RMSEA < 0.001$ ,  $AIC = 48.6$ ,  $r^2 = 0.37$ ,  $n = 590$ ).  $CO_2$  concentration and soil type, the latter represented as  $\log_e(\% \text{ clay})$  content), are independent variables. The growing season mean of soil water potential to 50 cm depth ( $\Psi_s$ ), the fractional contribution of the tallgrass *Sorghastrum* to community productivity ( $fSorg$ ) and  $fSorg$  during the previous year ( $fSorg_p$ ) are dependent variables.  $CO_2$  and soil type co-vary, as indicated by the curved line, because the physical placement of monoliths of a given soil type along the Lysimeter  $CO_2$  Gradient (LYCOG) facility was correlated with  $CO_2$  treatment. Standardized coefficients are listed beside the paths linking the variables. Solid lines represent positive correlations. Dashed lines represent negative correlations. Numbers displayed above the boxes for independent variables are estimated variances for [ $CO_2$ ]/100 or  $\log_e(\% \text{ clay})$ . Numbers displayed adjacent to the ovals are residual errors for dependent variables (e.g.  $e_{fSorg}$ ).

preceded and followed 2011 (Fig. S2). The  $CO_2$  effect on *Sorghastrum* abundance declined on all soils following 2011, possibly because increased evaporative demand disadvantaged this relatively drought-sensitive species (Swemmer *et al.*, 2006). Community change plays an important role in ecosystem response to global change drivers (Langley & Hungate, 2014), as it did in this experiment.

The predominant direct effect of  $CO_2$  on ANPP is probably explained by increased photosynthetic WUE (assimilation rate/transpiration rate) (Fay *et al.*, 2012). Evapotranspiration rates were similar across the  $CO_2$  gradient for prairie assemblages growing on a given soil type (Polley *et al.*, 2014); consequently, the  $CO_2$ -caused increase in WUE probably arose from increased canopy-level C assimilation rates. The genesis of the direct effect of soil percentage clay content on ANPP is difficult to discern from the available data, but soil effects were not explained by soil differences in water or N availability. Other mechanisms which may have contributed to a soil effect on productivity include soil differences in aeration, resistance to root penetration (Masle & Passioura, 1987; Wallace, 1987; Schrama *et al.*, 2013) and

limitation of ANPP by elements other than N (Niklaus & Körner, 2004; Polley *et al.*, 2016).

$CO_2$  and soil effects on ANPP were additive in this study, as initial soil differences in species succession diminished over time. Additive effects of climate change factors on diversity and productivity were also observed during the initial years of the Jasper Ridge Global Change Experiment (Shaw *et al.*, 2002; Zavaleta *et al.*, 2003). The more typical finding is that  $CO_2$  effects on ANPP are conditioned by other productivity-regulating variables (e.g. Reich & Hobbie, 2013; Mueller *et al.*, 2016; Zhu *et al.*, 2016). Whether additive or interactive effects on ANPP prevail across soil types will probably depend on the extent to which growth-limiting resources vary among soils. The greater the range in resource levels among soils, the greater the likelihood that the ANPP- $CO_2$  response will differ among soils. Under the conditions of this experiment, however,  $CO_2$  enrichment had a large and positive influence on grassland productivity that was expressed similarly across soil types. The initial ANPP- $CO_2$  response differed among soils because  $CO_2$  effects on  $\Psi_s$  and *Sorghastrum* abundance differed among soils. Soil-related differences in  $CO_2$  effects weakened during the

latter half of the experiment and did not influence across-soil responses over the decade-long study. Our results demonstrate a role of species differences and change in mediating soil effects on the ANPP–CO<sub>2</sub> response, but imply that CO<sub>2</sub> effects on productivity can be estimated across soils of different physical properties in some situations.

## Acknowledgements

Chris Kolodziejczyk operated the CO<sub>2</sub> chambers. Field and laboratory assistance from Katherine Jones, Chris Kolodziejczyk, Alicia Naranjo, Kyle Tiner and numerous researchers and students was invaluable. The mention of trade names or commercial products does not imply endorsement by the US Department of Agriculture (USDA). USDA is an equal opportunity provider and employer. The authors have no conflicts of interest to declare.

## Author contributions

HWP and PAF supervised LYCOG operation and wrote the initial draft of the manuscript. All authors, including MJA, HPC, AEG, RAG, RBJ, VLJ, ARK and LGR, contributed to data collection, processing and summarization, and aided in writing the final version of the manuscript.

## ORCID

Michael J. Aspinwall  <http://orcid.org/0000-0003-0199-2972>  
 Philip A. Fay  <http://orcid.org/0000-0002-8291-6316>  
 Richard A. Gill  <http://orcid.org/0000-0001-8981-0621>  
 Robert B. Jackson  <http://orcid.org/0000-0001-8846-7147>  
 Virginia L. Jin  <http://orcid.org/0000-0003-4146-8497>  
 Albina R. Khasanova  <http://orcid.org/0000-0001-6652-012X>  
 H. Wayne Polley  <http://orcid.org/0000-0002-1197-8800>  
 Lara G. Reichmann  <http://orcid.org/0000-0002-9737-7300>

## References

- Diggs GM Jr, Lipscomb BL, O'Kennon RJ. 1999. *Shinner's and Mahler's illustrated flora of North Central Texas*. Fort Worth, TX, USA: Botanical Research Institute of Texas Inc.
- Dukes JS, Chiariello NR, Cleland EE, Moore LA, Shaw MR, Thayer S, Tobeck T, Mooney HA, Field CB. 2005. Responses of grassland production to single and multiple global environmental changes. *PLoS Biology* 3: 1829–1837.
- Epstein HE, Lauenroth WK, Burke IC, Coffin DP. 1997. Productivity patterns of C<sub>3</sub> and C<sub>4</sub> functional types in the U.S. Great Plains. *Ecology* 78: 722–731.
- Epstein HE, Lauenroth WK, Burke IC, Coffin DP. 1998. Regional productivities of plant species in the Great Plains of the United States. *Plant Ecology* 134: 173–195.
- Fatichi S, Leuzinger S, Paschalis A, Langley JA, Barraclough AD, Hovenden MJ. 2016. Partitioning direct and indirect effects reveals the response of water-limited ecosystems to elevated CO<sub>2</sub>. *Proceedings of the National Academy of Sciences, USA* 113: 12757–12762.
- Fay PA, Jin VL, Way DA, Potter KN, Gill RA, Jackson RB, Polley HW. 2012. Soil-mediated effects of subambient to increased carbon dioxide on grassland productivity. *Nature Climate Change* 2: 742–746.
- Fay PA, Kelley AM, Procter AC, Hui D, Jin VL, Jackson RB, Johnson HB, Polley HW. 2009. Primary productivity and water balance of grassland vegetation on three soils in a continuous CO<sub>2</sub> gradient: initial results from the lysimeter CO<sub>2</sub> gradient experiment. *Ecosystem* 12: 699–714.
- Fay PA, Newingham BA, Polley HW, Morgan JA, LeCain DR, Nowak RS, Smith SD. 2015a. Dominant plant taxa predict plant productivity responses to CO<sub>2</sub> enrichment across precipitation and soil gradients. *AoB Plants* 7: plv027.
- Fay PA, Reichmann LG, Aspinwall MJ, Khasanova AR, Polley HW. 2015b. A CO<sub>2</sub> concentration gradient facility for testing CO<sub>2</sub> enrichment and soil effects on grassland ecosystem function. *Journal of Visualized Experiments* 105: e53151.
- García-Palacios P, Maestre FT, Bardgett RD, de Kroon H. 2012. Plant responses to soil heterogeneity and global environmental change. *Journal of Ecology* 100: 1303–1314.
- Grace JB. 2006. *Structural equation modeling and natural systems*. Cambridge, UK: Cambridge University Press.
- Hoerling M, Kumar A, Dole R, Nielsen-Gammon JW, Eischeid J, Perlwitz J, Quan X-W, Zhang T, Pegion P, Chen M. 2013. Anatomy of an extreme event. *Journal of Climate* 26: 2811–2832.
- Hovenden MJ, Newton PCD, Willis KE. 2014. Seasonal not annual rainfall determines grassland biomass response to carbon dioxide. *Nature* 511: 583–586.
- Hungate BA, Stiling PD, Dijkstra P, Johnson DW, Ketterer ME, Hymus GJ, Hinkle CR, Drake BG. 2004. CO<sub>2</sub> elicits long-term decline in nitrogen fixation. *Science* 304: 1291.
- Johnson HB, Polley HW, Whitis RP. 2000. Elongated chambers for field studies across atmospheric CO<sub>2</sub> gradients. *Functional Ecology* 14: 388–396.
- Langley JA, Hungate BA. 2014. Plant community feedbacks and long-term ecosystem responses to multi-factored global change. *AoB Plants* 6: plu035.
- Langley JA, Megonigal JP. 2010. Ecosystem response to elevated CO<sub>2</sub> levels limited by nitrogen-induced plant species shift. *Nature* 466: 96–99.
- Le Quéré C, Andrew RM, Friedlingstein P, Stich S, Pongratz J, Manning AC, Korsbakken JI, Petters GP, Canadell JG, Jackson RB *et al.* 2018. Global carbon budget 2017. *Earth System Science Data* 10: 405–448.
- Luo Y, Su B, Currie WS, Dukes JS, Finzi A, Hartwig U, Hungate B, McMurtrie RE, Oren R, Parton WJ *et al.* 2004. Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *BioScience* 54: 731–739.
- Masle J, Passioura JB. 1987. The effect of soil strength on the growth of young wheat plants. *Australian Journal of Plant Physiology* 14: 643–656.
- Morgan JA, Pataki DE, Körner C, Clark H, Del Grosso SJ, Grünzweig M, Knapp AK, Mosier AR, Newton PCD, Niklaus PA *et al.* 2004. Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO<sub>2</sub>. *Oecologia* 140: 11–25.
- Mueller KE, Blumenthal DM, Pendall E, Carrillo Y, Dijkstra FA, Williams DG, Follett RF, Morgan JA. 2016. Impacts of warming and elevated CO<sub>2</sub> on a semi-arid grassland are non-additive, shift with precipitation, and reverse over time. *Ecology Letters* 19: 956–966.
- Newton PCD, Lieffering M, Parsons AJ, Brock SC, Theobald PW, Hunt CL, Luo D, Hovenden MJ. 2014. Selective grazing modifies previously anticipated responses of plant community composition to elevated CO<sub>2</sub> in a temperate grassland. *Global Change Biology* 20: 158–169.
- Niklaus PA, Körner Ch. 2004. Synthesis of a six-year study of calcareous grassland responses to *in situ* CO<sub>2</sub> enrichment. *Ecological Monographs* 74: 491–511.
- Niklaus PA, Leadley PW, Schmid B, Körner Ch. 2001. A long-term field study on biodiversity × elevated CO<sub>2</sub> interactions in grassland. *Ecological Monographs* 71: 341–356.
- Nowak RS, Ellsworth DS, Smith SD. 2004. Functional responses of plants to elevated atmospheric CO<sub>2</sub> – do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist* 162: 253–280.
- Owensby CE, Auen LM, Coyne PI. 1994. Biomass production in a nitrogen-fertilized, tallgrass prairie ecosystem exposed to ambient and elevated levels of CO<sub>2</sub>. *Plant and Soil* 165: 105–113.
- Owensby CE, Ham JM, Knapp AK, Auen LM. 1999. Biomass production and species composition change in a tallgrass prairie ecosystem after long-term exposure to elevated atmospheric CO<sub>2</sub>. *Global Change Biology* 5: 497–506.
- Polley HW, Collins HP, Reichmann LG, Fay PA. 2016. CO<sub>2</sub> and soil water potential as regulators of the growth and N fraction derived from fixation of a legume in tallgrass prairie communities. *Plant and Soil* 409: 361–370.

- Polley HW, Derner JD, Jackson RB, Wilsey BJ, Fay PA. 2014. Impacts of climate change drivers on C<sub>4</sub> grassland productivity: scaling driver effects through the plant community. *Journal of Experimental Botany* 65: 3415–3424.
- Polley HW, Jin VL, Fay PA. 2012. Feedback from plant species change amplifies CO<sub>2</sub> enhancement of grassland productivity. *Global Change Biology* 18: 2813–2823.
- Polley HW, Johnson HB, Fay PA, Sanabria J. 2008. Initial response of evapotranspiration from tallgrass prairie vegetation to CO<sub>2</sub> at subambient to elevated concentrations. *Functional Ecology* 22: 163–171.
- Polley HW, Morgan JA, Fay PA. 2011. Application of a conceptual framework to interpret variability in rangeland responses to atmospheric CO<sub>2</sub> enrichment. *Journal of Agricultural Science* 149: 1–14.
- Polley HW, Wilsey BJ, Derner JD. 2007. Dominant species constrain effects of species diversity on temporal variability in biomass production of tallgrass prairie. *Oikos* 116: 2044–2052.
- Reich PB, Hobbie SE. 2013. Decade-long soil nitrogen constraint on the CO<sub>2</sub> fertilization of plant biomass. *Nature Climate Change* 3: 278–282.
- Reich PB, Hobbie SE, Lee TD, Pastore MA. 2018. Unexpected reversal of C<sub>3</sub> versus C<sub>4</sub> grass response to elevated CO<sub>2</sub> during a 20-year field experiment. *Science* 360: 317–320.
- Sala OE, Lauenroth WK, McNaughton SJ, Rusch G, Zhang X. 1996. Biodiversity and ecosystem functioning in grasslands. In: Mooney HA, Cushman JH, Medina E, Sala OE, Schulze ED, eds. *Functional roles of biodiversity: a global perspective*. Chichester, UK: Wiley, 129–149.
- Sala OE, Parton WJ, Joyce LA, Lauenroth WK. 1988. Primary production of the central grassland region of the United States. *Ecology* 69: 40–45.
- Schrama MJJ, Cordlandwehr V, Visser EJW, Elzenga TM, de Vries Y, Bakker JP. 2013. Grassland cutting regimes affect soil properties, and consequently vegetation composition and belowground plant traits. *Plant and Soil* 366: 401–413.
- Shaw MR, Zavaleta ES, Chiariello NR, Cleland EE, Mooney HA, Field CB. 2002. Grassland responses to global environmental changes suppressed by elevated CO<sub>2</sub>. *Science* 298: 1987–1990.
- Shipley B. 2000. *Cause and correlation in biology: a user's guide to path analysis, structural equations and causal inference*. Cambridge, UK: Cambridge University Press.
- Swemmer AM, Knapp AK, Smith MD. 2006. Growth responses of two dominant C<sub>4</sub> grass species to altered water availability. *International Journal of Plant Sciences* 167: 1001–1010.
- Tor-Ngern P, Oren R, Oishi AC, Uebelherr JM, Palmroth S, Tarvainen L, Ottosson-Löfvenius M, Linder S, Domec J-C, Nasholm T. 2017. Ecophysiological variation of pine forests: synthesis of new and published results. *Ecological Applications* 27: 118–133.
- Wallace LL. 1987. Effects of clipping and soil compaction on growth, morphology and mycorrhizal colonization of *Schizachyrium scoparium*, a C<sub>4</sub> bunchgrass. *Oecologia* 72: 423–428.
- Zavaleta ES, Shaw MR, Chiariello NR, Mooney HA, Field CB. 2003. Additive effects of simulated climate changes, elevated CO<sub>2</sub>, and nitrogen deposition on grassland diversity. *Proceedings of the National Academy of Sciences, USA* 100: 7650–7654.
- Zhu K, Chiariello NR, Tobeck T, Fukami T, Field CB. 2016. Nonlinear, interacting responses to climate limit grassland production under global change. *Proceedings of the National Academy of Sciences, USA* 113: 10589–10594.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article:

**Fig. S1** Aerial view of the Lysimeter CO<sub>2</sub> Gradient (LYCOG) facility.

**Fig. S2** Yearly irrigation and temperature in the CO<sub>2</sub> gradient experiment.

**Fig. S3** Structural equation model of variables affecting the annual aboveground net primary productivity (log<sub>e</sub>(ANPP)) of grassland communities during the first 5 yr of CO<sub>2</sub> treatment, 2006–2010.

**Fig. S4** Structural equation model of variables affecting the annual aboveground net primary productivity (log<sub>e</sub>(ANPP)) of grassland communities during the final 5 yr of CO<sub>2</sub> treatment, 2011–2015.

**Table S1** Linear, mixed model tests of the responses of productivity, species abundances, and soil water and N to independent variables and variable interactions over a decade-long CO<sub>2</sub> enrichment experiment.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.