

Plant community change mediates the response of foliar $\delta^{15}\text{N}$ to CO_2 enrichment in mesic grasslands

**H. Wayne Polley, Justin D. Derner,
Robert B. Jackson, Richard A. Gill,
Andrew C. Procter & Philip A. Fay**

Oecologia


ISSN 0029-8549

Volume 178

Number 2

Oecologia (2015) 178:591-601

DOI 10.1007/s00442-015-3221-x



Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg (outside the USA). This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Plant community change mediates the response of foliar $\delta^{15}\text{N}$ to CO_2 enrichment in mesic grasslands

H. Wayne Polley · Justin D. Derner ·
Robert B. Jackson · Richard A. Gill ·
Andrew C. Procter · Philip A. Fay

Received: 5 June 2014 / Accepted: 3 January 2015 / Published online: 22 January 2015
© Springer-Verlag Berlin Heidelberg (outside the USA) 2015

Abstract Rising atmospheric CO_2 concentration may change the isotopic signature of plant N by altering plant and microbial processes involved in the N cycle. CO_2 may increase leaf $\delta^{15}\text{N}$ by increasing plant community productivity, C input to soil, and, ultimately, microbial mineralization of old, ^{15}N -enriched organic matter. We predicted that CO_2 would increase aboveground productivity (ANPP; g biomass m^{-2}) and foliar $\delta^{15}\text{N}$ values of two grassland communities in Texas, USA: (1) a pasture dominated by a C_4 exotic grass, and (2) assemblages of tallgrass prairie species, the latter grown on clay, sandy loam, and silty clay soils. Grasslands were exposed in separate experiments

to a pre-industrial to elevated CO_2 gradient for 4 years. CO_2 stimulated ANPP of pasture and of prairie assemblages on each of the three soils, but increased leaf $\delta^{15}\text{N}$ only for prairie plants on a silty clay. $\delta^{15}\text{N}$ increased linearly as mineral-associated soil C declined on the silty clay. Mineral-associated C declined as ANPP increased. Structural equation modeling indicated that CO_2 increased ANPP partly by favoring a tallgrass (*Sorghastrum nutans*) over a mid-grass species (*Bouteloua curtipendula*). CO_2 may have increased foliar $\delta^{15}\text{N}$ on the silty clay by reducing fractionation during N uptake and assimilation. However, we interpret the soil-specific, $\delta^{15}\text{N}$ – CO_2 response as resulting from increased ANPP that stimulated mineralization from recalcitrant organic matter. By contrast, CO_2 favored a forb species (*Solanum dimidiatum*) with higher $\delta^{15}\text{N}$ than the dominant grass (*Bothriochloa ischaemum*) in pasture. CO_2 enrichment changed grassland $\delta^{15}\text{N}$ by shifting species relative abundances.

Communicated by Jim Ehleringer.

H. W. Polley (✉) · P. A. Fay
Grassland, Soil and Water Research Laboratory, USDA-
Agricultural Research Service, Temple, TX 76502, USA
e-mail: wayne.polley@ars.usda.gov

J. D. Derner
High Plains Grasslands Research Station, USDA-Agricultural
Research Service, Cheyenne, WY 82009, USA

R. B. Jackson
Nicholas School of the Environment, Duke University,
Durham, NC 27708, USA

R. B. Jackson
School of Earth Sciences, Stanford University,
Stanford, CA 94305, USA

R. A. Gill
Department of Biology, Brigham Young University,
Provo, UT 84602, USA

A. C. Procter
US Environmental Protection Agency, Research Triangle Park,
Durham, NC 27709, USA

Keywords Isotope · Plant productivity · Soil carbon ·
Soil type · Tallgrass prairie

Introduction

Atmospheric CO_2 enrichment commonly leads to changes in the N cycle because of the tight coupling that exists between organic forms of C and N (Luo et al. 2004; De Graaff et al. 2006; Gill et al. 2006). Plant ^{15}N natural abundances provide an integrated metric of how N-containing pools in soils and plants are transformed, accessed, and mixed (Robinson 2001) and thus can be used to interpret net effects of CO_2 on the N cycle. Plant $\delta^{15}\text{N}$ values, reflective of the ratio of ^{15}N – ^{14}N , have been found to decrease, increase, or not change at elevated CO_2 (Billings et al.




Fig. 1 Directional impacts of selected soil and plant processes on the $\delta^{15}\text{N}$ of plants (*Plant*) relative to an initial $\delta^{15}\text{N}$ value of the N pool in soil solution (*Soil Nitrogen*). Fractionating processes associated with denitrification or plant assimilation of NH_4^+ or NO_3^- exert their largest effect on plant $\delta^{15}\text{N}$ when the ratio of the supply rate or concentration of soil inorganic N is high relative to plant demand for N (ratio > 1). Conversely, $\delta^{15}\text{N}$ values are more influenced by plant exploitation of alternative N sources (e.g., mineralization of recalcitrant SOM; increased acquisition of N from deep soil layers) or mechanisms of N uptake (e.g., via mycorrhizae, *M+*) when N supply approaches plant demand (ratio ≈ 1)

2002; Bassirirad et al. 2003; Stock and Evans 2006; Garten et al. 2011), testament to both the diversity of N cycle processes potentially influenced by CO_2 and the range of plant and ecosystem responses to CO_2 .

CO_2 -caused changes in plant $\delta^{15}\text{N}$ signatures result from changes in soil- or plant-mediated processes that affect or are influenced by isotopic fractionation. CO_2 effects on plant $\delta^{15}\text{N}$ are expected to vary as a function of the supply rate of inorganic N in soil solution relative to the demand for organic N by plants (Fig. 1; Evans 2001; Robinson 2001; Kalcsits et al. 2014). Fractionation may occur during N transformations in soil and plant uptake, assimilation, and turnover/export of N (Evans 2001; Robinson 2001; Soper et al. 2014). Fractionation is expected to be greatest when N supply exceeds demand. For example, discrimination during nitrate and ammonium assimilation in roots is thought to occur when plant pools of inorganic N are only partially consumed and inorganic N is lost from roots to the surrounding substrate (Evans 2001; Kalcsits et al. 2014). Assimilated plant N will be depleted in ^{15}N relative to source N when N supply exceeds plant demand and fractionation is expressed, but organic and inorganic plant N will not differ in $\delta^{15}\text{N}$ signature if the entire inorganic pool in plants is assimilated (Kalcsits et al. 2014). An excess in N supply relative to plant demand could also lead to an increase in the $\delta^{15}\text{N}$ signature of inorganic N in soil solution, likely without changing the $\delta^{15}\text{N}$ signature of the larger N pool in bulk soil. Solution N is enriched in ^{15}N by

losses of ^{15}N -depleted N molecules to leaching or denitrification (Robinson and Conroy 1999; Garten et al. 2011).

CO_2 is usually assumed to influence plant $\delta^{15}\text{N}$ by increasing plant demand for N relative to supply, resulting in a change in the isotopic signature of plant-available soil N. CO_2 could change the $\delta^{15}\text{N}$ of the plant-available N pool by causing the addition or loss of isotopically distinct N or permitting plants to access a previously under-exploited pool of isotopically distinct N (Robinson 2001). The progressive N limitation (PNL) hypothesis holds that CO_2 enrichment will intensify N limitation of plant production by increasing C input to soil and N sequestration in soil organic matter (SOM) or long-lived plant tissues (Luo et al. 2004). This tightening of the N cycle is anticipated to reduce N losses. A decline in denitrification would reduce the $\delta^{15}\text{N}$ of soil NO_3^- (Robinson and Conroy 1999), for example, whereas increasing denitrification is thought to contribute to the positive correlation that is often observed between $\delta^{15}\text{N}$ values and foliar N concentration or soil N availability or mineralization rate (Kahmen et al. 2008; Craine et al. 2009). Conversely, CO_2 enrichment could increase plant $\delta^{15}\text{N}$ in systems in which N losses are minimal by increasing soil microbial activity (Billings et al. 2002) and microbial immobilization of N or mineralization of old, recalcitrant organic matter (Johnson DW et al. 2000; Gill et al. 2002; Dijkstra et al. 2010), the latter of which is enriched in ^{15}N (Tiessen et al. 1984). The PNL theory also holds that effects of a CO_2 -caused decline in N availability to plants can initially be overcome by a transfer of N from low C:N ratio fractions, such as recalcitrant SOM, to plants with higher C:N ratios. Additionally, CO_2 enrichment could alter plant reliance on mycorrhizal fungi for N uptake (Bassirirad et al. 2003). Nitrogen derived from mycorrhizal associates is depleted in ^{15}N compared to source N (Evans 2001; Hobbie and Hobbie 2008). Finally, plants may exhibit no change in $\delta^{15}\text{N}$ values, particularly when N is strongly limiting. Fractionation is reduced when plant demand approaches N supply rates (Robinson 2001; Billings et al. 2002).

Soil properties mediate several of the processes that determine how N containing pools are transformed, accessed, and mixed, and thus may affect how plant $\delta^{15}\text{N}$ values respond to CO_2 . Soil differences in texture and related hydrological properties potentially influence organic pools of N (Fay et al. 2009) and N mineralization and denitrification rates (Bechtold and Naiman 2006; Gu and Riley 2010), all of which may affect the plant $\delta^{15}\text{N}$ response to CO_2 .

Several of the processes that affect $\delta^{15}\text{N}$ values of plants or plant communities may be influenced indirectly by CO_2 -caused change in the composition or relative abundances of species in plant communities (community change). We consider effects of CO_2 -caused community change on

abundance-weighted (community) values of leaf $\delta^{15}\text{N}$ to be species-specific when the species involved differ in $\delta^{15}\text{N}$ values. CO_2 could alter community $\delta^{15}\text{N}$ by differentially favoring plant species that differ in the form or source of N accessed or within-plant fractionation. Legumes often have leaf $\delta^{15}\text{N}$ signatures near zero because of their reliance on symbiotic N_2 fixation (Robinson 2001; Hungate et al. 2004). Other species may differ in $\delta^{15}\text{N}$ signatures because of differing preferences for NH_4^+ compared to NO_3^- (Robinson 2001; Kahmen et al. 2008; Wang and Macko 2011) or rooting depth (Handley and Scrimgeour 1997), for example. Alternatively, increased CO_2 could shift the $\delta^{15}\text{N}$ signatures of species by a similar magnitude and direction by favoring species that disproportionately influence soil C inputs, microbial activity, and, ultimately, the $\delta^{15}\text{N}$ signature of N in soil solution. CO_2 effects on community $\delta^{15}\text{N}$ values are non-species-specific when the magnitude and direction of change in $\delta^{15}\text{N}$ are similar among species in the community.

We use data from two experiments on grasslands to determine how leaf $\delta^{15}\text{N}$ values of dominant perennial species responded to a pre-industrial to elevated gradient in CO_2 concentration and to assess possible explanations for $\delta^{15}\text{N}$ change. The first grassland is a formerly-grazed pasture dominated by an exotic C_4 grass. The second grassland consists of assemblages of native tallgrass prairie species grown on each of three soil types. Both grasslands were exposed to the CO_2 gradient for 4 years during separate experiments. CO_2 enrichment increased ANPP (g biomass m^{-2}) and changed community composition in both pasture (Polley et al. 2003) and prairie assemblages (Fay et al. 2012; Polley et al. 2012a). The increase in ANPP and shift in communities was accompanied by increased demand for N in aboveground plant tissues (Gill et al. 2006; Polley et al. 2011). Likely as a consequence of heightened plant demand for N, we observed changes in N mineralization rates, the C:N ratio of SOM, soil fungal communities, and activities of microbially-derived extracellular enzymes involved in C and N cycling in soil, and found evidence for a net transfer of N from recalcitrant SOM with low C:N ratios to plants with larger C:N ratios (Gill et al. 2002, 2006; Kelley et al. 2011; Procter et al. 2014). CO_2 enrichment has repeatedly been shown to alter soil microbial dynamics and N cycling and, in some ecosystems, to increase N transfer from soils to plants (King et al. 2004; Dijkstra et al. 2010) apparently by stimulating ANPP and C input to soil. Yet, there remains little direct or even correlative evidence that plant N dynamics are influenced by CO_2 -caused changes in soil microbial processes. We predicted that CO_2 enrichment would increase foliar $\delta^{15}\text{N}$ values in pasture and prairie assemblages, consistent with evidence that CO_2 stimulated both ANPP (Polley et al. 2003, 2012b; Fay et al. 2012) and mineralization of

recalcitrant SOM (Gill et al. 2002). Mineralization of recalcitrant SOM would be expected to increase $\delta^{15}\text{N}$ values of plant available N (Johnson DW et al. 2000; Billings et al. 2002). Greater productivity may increase C input to soil and thereby stimulate microbes to decompose relatively old SOM (Gill et al. 2002; Dijkstra et al. 2010). Consequently, we anticipated that foliar $\delta^{15}\text{N}$ values would increase with increased ANPP. The magnitude of the ANPP increase potentially is influenced by soil properties (Fay et al. 2012) and community dynamics (Polley et al. 2014). We predicted that CO_2 enrichment would shift the $\delta^{15}\text{N}$ signatures of species by a similar magnitude and direction by favoring species that augment the ANPP- CO_2 response and, ultimately, soil C input and microbial activity. An alternative prediction is that CO_2 will shift community $\delta^{15}\text{N}$ by altering relative abundances of species that differ in $\delta^{15}\text{N}$ values.

Materials and methods

Experimental design

We report results from two experiments in which elongated field chambers were used to expose vegetation to a continuous gradient in CO_2 spanning pre-industrial to elevated concentrations. In 1997–2000, we studied CO_2 effects on previously grazed C_3/C_4 grassland (hereafter, pasture) using the Prairie CO_2 Gradient (PCG) facility (Johnson HB et al. 2000). In 2006–2009, we evaluated CO_2 effects on assemblages of tallgrass prairie species grown in soils of three types using the Lysimeter CO_2 gradient (LYCOG) facility (Polley et al. 2008; Fay et al. 2009). Both facilities were located in central Texas USA (31°05'N, 97°20'W) and consisted of two transparent and tunnel-shaped chambers, aligned parallel along a north–south axis. Each chamber was divided into ten consecutive compartments each 5 m long and 1.0 m (PCG) or 1.2 m (LYCOG) wide and tall. Chambered vegetation was enclosed in a transparent polyethylene film. Photosynthesis by enclosed vegetation progressively depleted the CO_2 concentration in air as it was moved by blowers toward the air outlet of each chamber to create daytime CO_2 gradients of 560 (PCG) or 500 $\mu\text{L L}^{-1}$ (LYCOG) to 370 $\mu\text{L L}^{-1}$ (elevated chamber) and 370 $\mu\text{L L}^{-1}$ to 200 (PCG) or 250 $\mu\text{L L}^{-1}$ (LYCOG) (subambient chamber). Night-time CO_2 concentrations were regulated at 130–150 $\mu\text{L L}^{-1}$ above daytime values along each chamber. Air temperature and vapor pressure deficit were regulated near ambient values by cooling and dehumidifying air at 5-m intervals along chambers. CO_2 treatments were maintained each growing season from March/April through mid-November.

The earlier PCG facility was constructed on grassland dominated by the C_4 perennial grass *Bothriochloa*

ischaemum (L.) Keng and the C₃ perennial forbs *Solanum dimidiatum* Raf., *Ratibida columnaris* (Sims) D. Don, and *Solidago canadensis* L. (hereafter referenced by genus; Polley et al. 2003). The site had been grazed for at least 50 years prior to construction. The soil was a silty clay. Soil beneath chambers was separated from surrounding soil to a depth of 0.9 m with a rubber-coated fabric. The LYCOG facility was constructed atop 5-m-long × 1.2-m-wide × 1.6-m-deep steel containers that were buried to 1.2 m depth. Two intact soil monoliths (each 1 × 1 × 1.5 m deep) of each of two soil types were placed into each of the 20 5-m-long containers. Three soil types of contrasting physical and hydrological properties were represented: (1) silty clay on which the PCG facility was constructed, (2) clay, and (3) sandy loam (Fay et al. 2009). Two monoliths of each of two soil types were randomly placed in each 5-m-long compartment. Four perennial C₄ grass species and three perennial C₃ forb species, all characteristic of tallgrass prairie in central Texas, were transplanted into 60 of the total of 80 monoliths in June 2003, 3 years prior to CO₂ treatment (Polley et al. 2008; Fay et al. 2009). Eventual dominants included the C₄ grasses *Bouteloua curtipendula* (Michx.) Torr., and *Sorghastrum nutans* (L.) Nash and the forb species *Solidago canadensis* (hereafter referenced by genus).

Irrigation equivalent to rainfall was applied to grassland in the PCG facility on the day following precipitation (Polley et al. 2002). Each monolith in the later LYCOG facility was irrigated twice weekly during each growing season. Irrigation was applied to simulate the seasonal distribution and average of growing season precipitation in central Texas (560 mm; Polley et al. 2011).

Vegetation and soil sampling and data analysis

Soils and upper canopy leaves (or leaf blades) were sampled from along the PCG and LYCOG experiments for N isotope measurements. Soil cores (4.2 cm diameter, 53 cm depth) were collected in December 2000 at 1-m distance from both the air entrance and exit of each 5-m compartment of the PCG. Soils from the two cores per compartment were composited over 0–8, 8–23, 23–37, and 37–53 cm depth increments. We collected fully expanded leaves of *Solanum* in June 2000 and of *Bothriochloa* in June 1997 and 2000, the first and fourth years of the PCG experiment, and leaves of the forb *Solidago* and two grasses, *Bouteloua* and *Sorghastrum*, in June 2005 and 2009, the year prior to CO₂ treatment and fourth year of the LYCOG experiment. Grasslands in central Texas usually approach maximal physiological activity during June (Mielnick et al. 2001). Leaves were collected from multiple individuals of each of the target species, when possible, and composited by species for each 1-m² area sampled along the CO₂ gradient

(PCG) or for each monolith (LYCOG). ANPP of each species was determined by clipping vegetation to 5-cm height at the end of each growing season.

The N isotope composition of soil samples and whole leaves was determined by mass spectrometry and expressed as δ¹⁵N, ‰ (parts per thousand) ¹⁵N relative to air (PCG–Isotope Services, Los Alamos, NM, USA; LYCOG–University of California Davis Stable Isotope Facility, Davis, CA, USA). The N content of samples was measured during isotope analyses and expressed as [N], mg N g⁻¹ biomass. We measured the N isotope signature of top-soil samples (8–23 cm depth increment) for each 5-m compartment and of soil from three additional depth increments (0–8, 23–37, and 37–53 cm) for each of three compartments (at 550, 430, 360 μL L⁻¹ CO₂; PCG). We included an additional five leaf samples in each analysis of plant materials as internal standards. Data from these samples were used to standardize results from multiple measurement sessions for each experiment (PCG, LYCOG). Results for standards from any two measurement sessions were highly correlated ($r^2 > 0.97$). Pre-treatment to post-treatment changes in δ¹⁵N values during each experiment were calculated by subtracting δ¹⁵N measured on leaves collected in 1997 or 2005 from δ¹⁵N of leaves from 2000 or 2009 (PCG and LYCOG experiments, respectively).

LYCOG soils were sampled to 15-cm depth following the fourth season of CO₂ treatment (2009) to determine SOM fractions. Samples from the two monoliths of a given soil type in each 5-m chamber compartment were combined for physical fractionation following Gill (2007). SOM was fractionated into three components: coarse particulate organic matter (POM) C (>250 μm), fine POM C (53–250 μm), and mineral-associated C (<53 μm). Coarse POM represents recently-added organic C, whereas mineral-associated C is the most recalcitrant size fraction (Cambardella and Elliott 1992).

Bivariate regression analysis was used to determine relationships between variables measured during the PCG experiment and CO₂ concentration. Differences in mean values for the two species from the PCG experiment were analyzed with paired *t* tests. Spearman rank-order correlation analysis was used to compare trends in leaf N concentration, [N], between species pairs from the LYCOG experiment.

Data from the LYCOG experiment were analyzed using a mixed-model analysis of covariance (ANCOVA). Analyses were conducted with SAS (Littell et al. 2002). Soil type and species identity were considered fixed effects. The assignment of soil types to 5-m lengths of chambers was considered a random effect. CO₂ treatment and [N] of leaves were treated as covariates in the analysis.

Structural equation modeling (SEM) with observed variables (path modeling) was used to partition the net effect of CO₂ on change in δ¹⁵N, when significant, into a direct effect

and indirect effects mediated through changes in community ANPP and the fractional contribution of dominant species to community ANPP (Shipley 2000; Grace 2006). Bivariate relationships between modeled variables were linear. The SEM model was fit using IBM SPSS AMOS 21 software. The hypothesized relationship among variables in a SEM is considered to be consistent with data when the probability level of the statistical test (chi-squared statistic) is greater than the significance level ($P = 0.05$; Shipley 2000). Data were standardized by subtracting the mean and dividing by the standard deviation prior to analysis.

Results

PCG (pasture) experiment

ANPP

The ANPP–CO₂ response of pasture declined from the first to fourth years of treatment (1997–2000), but averaged 120 g m⁻² per 100 μL L⁻¹ increase in CO₂ over the first 3 years of the experiment (Polley et al. 2003). The 1997–2000 change in ANPP of the grass *Bothriochloa* was a negative linear function of CO₂ (Fig. 2a). ANPP of *Bothriochloa* increased from the first to fourth years of treatment at CO₂ levels <320 μL L⁻¹, but declined at higher CO₂. The 1997–2000 change in ANPP of the forb *Solanum* was not correlated with CO₂ ($P = 0.27$), but was greater on average over elevated than subambient concentrations (increase of 49 and 30 g biomass m⁻², respectively). The ratio of *Solanum* ANPP to *Bothriochloa* ANPP increased from a mean of 0.05 in 1997 to 0.18 and 0.30 at subambient and elevated CO₂, respectively, in 2000. Together, the two species contributed 60 and 30 % of community ANPP in the first and fourth years across CO₂ treatments.

δ¹⁵N

The δ¹⁵N of *Bothriochloa* leaves from both early in the initial treatment year of 1997 and final year of 2000 declined linearly at higher CO₂ (Fig. 3a). The slopes of δ¹⁵N–CO₂ regressions did not differ significantly between years for *Bothriochloa* ($P > 0.50$), but δ¹⁵N values were 1.27 ‰ lower (less positive) in 2000 than 1997. Thus, CO₂ did not affect the decline in leaf δ¹⁵N between 1997 and 2000. CO₂ also had no consistent effect on δ¹⁵N values of leaves of the forb *Solanum* in 2000. Leaf [N] and δ¹⁵N were not correlated for either *Bothriochloa* ($P = 0.21$) or *Solanum* ($P = 0.09$), but both [N] and δ¹⁵N were greater for the forb than the grass in the fourth treatment year (2000). Leaf [N] declined linearly at higher CO₂ for *Solanum* in 2000 and was greater on average by a factor of three for the forb




Fig. 2 Relationships between the change in ANPP (Δ ANPP) of the C₃ forb *Solanum* (square) and C₄ grass *Bothriochloa* (circle) over 4 years of the PCG experiment on pasture and CO₂ concentration (a) and between the N concentration, [N], of leaves of the forb and grass from year four of treatment and CO₂ (b). a Δ ANPP for *Bothriochloa* and b the [N]–CO₂ relationship for *Solanum* were fit with linear regressions (Δ ANPP = 502.61 – 1.56 × CO₂, adj. $r^2 = 0.44$, $P = 0.0008$, $n = 20$; [N] = 49.796 – 0.035 × CO₂, adj. $r^2 = 0.44$, $P = 0.0004$, $n = 22$). a Δ ANPP of *Solanum* and b leaf [N] of the *Bothriochloa* were not significantly related to CO₂ ($P = 0.27$ and 0.21, respectively)

(35.8 mg N g⁻¹ biomass) than grass (11.8 mg N g⁻¹ biomass; Fig. 2b). Leaf δ¹⁵N values averaged 5.58 and 1.66 ‰ across CO₂ treatments for *Solanum* and *Bothriochloa*, respectively (Fig. 3a). δ¹⁵N values of bulk soil increased with depth at all three positions sampled along the CO₂ gradient in 2000 (Fig. 3b).

LYCOG (tallgrass prairie) experiment

ANPP

CO₂ increased the 4-year average of ANPP of prairie communities by 100–110 g m⁻² per 100 μL L⁻¹ rise (Polley et al.




Fig. 3 Relationships between $\delta^{15}\text{N}$ values of bulk soil (as opposed to soil solution) or leaves from pasture and either CO_2 concentration (**a**) or soil depth (**b**) in the PCG experiment. **a** Soil (closed square) and leaves of the forb *Solanum* (open square) were collected in 2000. Leaf blades of the grass *Bothriochloa* were harvested during the first (closed circle, 1997) and fourth years of treatment (open circle, 2000). $\delta^{15}\text{N}$ – CO_2 relationships for bulk soil (8–23 cm depth increment; adj. $r^2 = 0.25$, $P = 0.02$, $n = 20$) and *Solanum* leaves (adj. $r^2 = 0.40$, $P = 0.003$, $n = 22$) were fit with linear and quadratic regression equations, respectively. Those for *Bothriochloa* from 1997 and 2000 were fit with parallel lines (slopes did not differ significantly, $P > 0.50$). **b** Relationships between $\delta^{15}\text{N}$ and soil depth (three locations along the CO_2 gradient) were fit with exponential equations (550 $\mu\text{L L}^{-1}$ CO_2 , adj. $r^2 = 0.99$, $P = 0.002$, $n = 4$; 430 and 360 $\mu\text{L L}^{-1}$ CO_2 combined; adj. $r^2 = 0.88$, $P = 0.0004$, $n = 8$)

2012b). The ANPP– CO_2 response was linear on the silty clay and sandy loam soils, but was strongly non-linear on the clay soil with little change in ANPP at $>390 \mu\text{L L}^{-1} \text{CO}_2$.

$\delta^{15}\text{N}$

The $\delta^{15}\text{N}$ values of leaves collected during the fourth year of the LYCOG experiment (2009) differed among species

Table 1 Results from a mixed-model analysis of covariance of species and soil effects on leaf $\delta^{15}\text{N}$ values and of slopes of regression relationships between $\delta^{15}\text{N}$ and both CO_2 and leaf [N] (LYCOG experiment)

Variable/unit	Mean/slope	P value
Species/ $\delta^{15}\text{N}$ (‰)		<0.0001
<i>Sorghastrum nutans</i>	1.3 (0.2) a	
<i>Bouteloua curtipendula</i>	0.0 (0.3) b	
<i>Solidago canadensis</i>	−0.5 (0.3) b	
Soil type		<0.0001
Clay	2.9 (0.3) a	
Sandy loam	−0.9 (0.3) b	
Silty clay	−1.3 (0.3) b	
Species X soil type		0.74
CO_2 (soil type)/‰ change $\delta^{15}\text{N}$ per 100 $\mu\text{L L}^{-1}$ increase in CO_2		0.001
Clay	–	0.35
Sandy loam	–	0.85
Silty clay	1.9	<0.0001
[N] (soil type)/‰ change $\delta^{15}\text{N}$ per 10 mg N g^{-1} biomass increase in [N]		<0.0001
Clay	3.2	0.0004
Sandy loam	–	0.15
Silty clay	3.9	<0.0001
CO_2 (species)		0.44
[N] (species)		0.04

P values indicate the significance of variables or variable interactions on leaf $\delta^{15}\text{N}$ and regression slopes. Means (SE) of $\delta^{15}\text{N}$ do not differ significantly among species ($n = 46$ –55) or soil types ($n = 40$ –60) if followed by the same letter

and plants grown on different soil types (Table 1). Leaf $\delta^{15}\text{N}$ was greater for the tallgrass species *Sorghastrum* than mid-grass *Bouteloua* and forb *Solidago* ($n = 46$, 51, and 55, respectively). Leaf $\delta^{15}\text{N}$ also was greater for plants grown on the clay than silty clay and sandy loam soils ($n = 52$, 60, and 40, respectively). Species effects on $\delta^{15}\text{N}$ did not depend on soil type, but the regression relationship between leaf $\delta^{15}\text{N}$ and CO_2 differed among soils. Leaf $\delta^{15}\text{N}$ was a significant linear function of CO_2 for the silty clay only (Fig. 4a). Regression relationships between $\delta^{15}\text{N}$ and CO_2 did not differ significantly among species (Table 1).

Leaf $\delta^{15}\text{N}$ was a significant and positive linear function of leaf [N] for the two clay soils (Table 1; Fig. 5). Leaf $\delta^{15}\text{N}$ increased by 3.2–3.9 ‰ for each 10 mg N g^{-1} biomass increase in [N]. Leaf [N] was not, however, correlated with CO_2 treatment for any soil type ($P = 0.63$, 0.30, 0.93 for silty clay, sandy loam, and clay soils, respectively). Leaf [N] was a negative function of species' fractional contributions to ANPP on the silty clay soil when analyzed using data for all species combined (not shown; adj. $r^2 = 0.18$, $P = 0.0005$, $n = 60$). There was no relationship between




Fig. 4 Relationships between leaf $\delta^{15}\text{N}$ values of perennial prairie species (*Bouteloua*, *Solidago*, *Sorghastrum*) grown on a silty clay soil and CO_2 concentration during the LYCOG experiment. Linear regressions between **a** $\delta^{15}\text{N}$ values of leaves and CO_2 treatment (year 4 of treatment; $\delta^{15}\text{N} = -8.346 + 0.019 \times \text{CO}_2$, adj. $r^2 = 0.28$, $P < 0.0001$, $n = 60$), and **b** between the 2005 and 2009 change in leaf $\delta^{15}\text{N}$ and CO_2 were derived from an analysis of covariance (change $\delta^{15}\text{N} = -5.845 + 0.014 \times \text{CO}_2$, adj. $r^2 = 0.23$, $P = 0.008$, $n = 37$)

leaf [N] and species' contributions to ANPP for the sandy loam ($P = 0.61$) or clay soils ($P = 0.50$). Together, CO_2 and leaf [N] explained 53 % of the variance in $\delta^{15}\text{N}$ values for the silty clay soil [$\delta^{15}\text{N} = -6.027 + 0.439 \times [\text{N}] + 0.018 \times r_{\text{CO}_2}$, where r_{CO_2} = variance in CO_2 concentration ($\mu\text{L L}^{-1}$) not explained by correlation with [N] (mg N g^{-1} biomass); $P < 0.0001$].

Change in $\delta^{15}\text{N}$

The pre-treatment to post-treatment (2005–2009) change in leaf $\delta^{15}\text{N}$ values generally was negative for prairie vegetation, indicating that $\delta^{15}\text{N}$ values declined during CO_2 treatment as observed for the PCG experiment (Fig. 4b). Analysis of covariance indicated that the temporal change in leaf




Fig. 5 Relationships between values of leaf $\delta^{15}\text{N}$ and N concentration, [N], for perennial prairie species grown on each of three soil types (clay, sandy loam, silty clay) during the LYCOG experiment. Linear regressions were derived from an analysis of covariance and were significant for the two clay soils (clay, $P = 0.0004$, $n = 52$; silty clay, $P < 0.0001$, $n = 60$)




Fig. 6 Structural equation model describing direct and indirect effects of CO_2 on the pre-treatment to post-treatment (4th year) change in leaf $\delta^{15}\text{N}$ values (change $\delta^{15}\text{N}$) of prairie species (*Bouteloua*, *Solidago*, *Sorghastrum*) grown on a silty clay soil during the LYCOG experiment ($n = 37$). We modeled indirect effects of CO_2 through the pre- to post-treatment change in both ANPP (ΔANPP) and the *Sorghastrum* fraction of community ANPP ($\Delta \text{Sorghastrum fraction}$) and a direct effect resulting from changes not linked to ANPP. The non-significant pathway is indicated by a *dashed line*. Standardized coefficients are listed beside each path

$\delta^{15}\text{N}$ did not differ among species ($P = 0.25$) or soil types ($P = 0.14$). Slopes of regression relationships between the change in leaf $\delta^{15}\text{N}$ and both CO_2 and leaf [N] did not differ among species ($P = 0.45$ and 0.36 , respectively). Regressions between change in leaf $\delta^{15}\text{N}$ and CO_2 did differ among soil types ($P = 0.02$).

The change in $\delta^{15}\text{N}$ was a significant linear function of CO_2 for the silty clay soil only (Fig. 4b). Leaf $\delta^{15}\text{N}$ declined from 2005 to 2009 at subambient CO_2 , but increased slightly at $>450 \mu\text{L L}^{-1} \text{CO}_2$. The positive effect of CO_2 on the change in $\delta^{15}\text{N}$ was mediated entirely through an increase in ANPP in a SEM that included possible CO_2

effects via mechanisms not linked to ANPP (Fig. 6). The more that ANPP increased from 2005 to 2009, the more the values of $\delta^{15}\text{N}$ increased. CO_2 enrichment increased ANPP both via a direct pathway and, indirectly, via community change associated with an increase in the *Sorghastrum* fraction. For the silty clay soil, community change increased $\delta^{15}\text{N}$ by increasing ANPP. We consider this impact of community change on foliar $\delta^{15}\text{N}$ to be non-species-specific, as it was expressed similarly among species (Fig. 4b).

Soil C fractions were strongly correlated with the 4-year change in ANPP along the CO_2 gradient in the silty clay soil (Fig. 7). Pools of the readily decomposable coarse POM fraction increased, whereas pools of recalcitrant, mineral-associated C decreased linearly as community ANPP increased. High values of change in leaf $\delta^{15}\text{N}$ were associated with reduced levels of mineral-associated C.

Would the quantity of N released during decomposition of mineral-associated SOM have been sufficient to supply N demands of the observed increase in ANPP at elevated CO_2 ? As estimated from regression (Fig. 7b), an increase in ANPP of $400 \text{ g biomass m}^{-2}$ was associated with an approximately 130 g C m^{-2} decline in the mineral-bound organic fraction in the silty clay soil (from 420 to 290 g C m^{-2}). Assuming a C:N ratio of 13–25 with extremes in C:N reflecting the measured ratio of total organic C to organic N (13) and the C:N of microaggregate POM in the silty clay (25; Gill et al. 2006), decomposition of 130 g m^{-2} of mineral-associated C may have released a maximum of $5\text{--}10 \text{ g N m}^{-2}$. CO_2 did not affect the [N] in aboveground tissues of dominant grass species harvested in June 2009, but reduced the [N] of grass assemblages by 6 % (from 9.3 to 8.7 mg N g^{-1} biomass) by increasing the relative abundance of *Sorghastrum*, the grass with the lowest [N] (Polley et al. 2011). At these plant N concentrations, a maximum of $3.5\text{--}3.7 \text{ g N m}^{-2}$ of additional N would have been required to fully supply demands of the $400 \text{ g biomass m}^{-2}$ increase in ANPP at elevated CO_2 . This additional N requirement represents approximately 35–75 % of the N potentially released during decomposition of mineral-associated SOM at elevated CO_2 .

The $\delta^{15}\text{N}$ value for newly-mineralized N can be estimated by assuming that mineralization of recalcitrant SOM fully supplied the N demands of increased ANPP. As much as 60 % of total ANPP at elevated CO_2 resulted from CO_2 stimulation of biomass production above the pre-treatment mean of $250 \text{ g biomass m}^{-2}$. The 2005–2009 change in leaf $\delta^{15}\text{N}$ was approximately -2 ‰ at subambient CO_2 (Fig. 4b). Temporal change in leaf $\delta^{15}\text{N}$ was approximately 0 ‰ in prairie species grown at $450 \mu\text{L L}^{-1}$ CO_2 . CO_2 stimulated ANPP by as much as $400 \text{ g biomass m}^{-2}$. Using a simple two-member isotope mixing model, we estimate that a change in $\delta^{15}\text{N}$ of approximately 1.3 ‰ from pre-treatment values would be required in the




Fig. 7 Relationships between organic C fractions in soil (coarse POM, **a**, or mineral-associated C, **b**) and the pre-treatment to post- CO_2 treatment change in ANPP of prairie vegetation (Δ ANPP) and between the pre- to post-treatment change in leaf $\delta^{15}\text{N}$ values (change leaf $\delta^{15}\text{N}$) of prairie species and C in the mineral-associated fraction (**c**). Data were derived from monoliths of the silty clay soil from along the CO_2 gradient in the LYCOG experiment. Data points represent mean values (Δ ANPP, change leaf $\delta^{15}\text{N}$) or results from composited samples (C fractions) from the two monoliths of silty clay soil per 5-m-long compartment along CO_2 chambers. Relationships were fit using linear regression (adj. $r^2 = 0.67, 0.47, 0.25$; $P = 0.0004, 0.006, 0.05$ for data in **a**, **b**, and **c**, respectively; $n = 13$)

400 g m^{-2} biomass added at elevated CO_2 to effect a net change in plant $\delta^{15}\text{N}$ of 0 ‰ as observed at $450 \mu\text{L L}^{-1}$ CO_2 (Fig. 4b).

Discussion

We predicted that foliar $\delta^{15}\text{N}$ values would (1) vary as a positive function of the amount by which CO_2 increased community ANPP, and (2) respond similarly among species. Our predictions were only partially supported. CO_2 enrichment increased ANPP of both pasture (PCG experiment; Polley et al. 2003) and prairie vegetation (LYCOG experiment; Fay et al. 2012; Polley et al. 2012b), but increased leaf $\delta^{15}\text{N}$ values to a similar extent for the three prairie species, and then only on a silty clay soil. The inconsistent response of $\delta^{15}\text{N}$ to CO_2 among soils resulted partly because CO_2 effects on prairie ANPP and species abundances differed among soil types. The silty clay soil on which CO_2 increased leaf $\delta^{15}\text{N}$ values was the least productive of the three soils on which prairie vegetation was grown, but displayed the most consistent ANPP– CO_2 responses among years (Fay et al. 2012; Polley et al. 2012b). Change in community composition, as reflected in a shift in dominance from a mid-grass to tall-grass species at elevated CO_2 , was the main driver of the large increase in ANPP on the silty clay. Increased ANPP, in turn, was linked with a decline in mineral-associated C in soil that may have contributed to greater plant reliance on ^{15}N -rich N derived from microbial decomposition of this recalcitrant pool of SOM. CO_2 enrichment did not consistently influence $\delta^{15}\text{N}$ values of either the dominant grass or forb in the PCG experiment, but increased the relative abundance of the forb species with higher $\delta^{15}\text{N}$ values than the dominant C_4 grass. Our results highlight a role of plant species (community) change in determining CO_2 effects on $\delta^{15}\text{N}$ values of grassland vegetation.

CO_2 altered community $\delta^{15}\text{N}$ values by favoring some species over others, but the mechanism involved in the feedback of community change on $\delta^{15}\text{N}$ differed between experiments. CO_2 effects were species-specific in the PCG experiment, as they resulted from change in the relative abundances of species that differed in $\delta^{15}\text{N}$ values. CO_2 enrichment favored a perennial forb with high leaf $\delta^{15}\text{N}$ values over the dominant grass with lower $\delta^{15}\text{N}$. Conversely, CO_2 effects on $\delta^{15}\text{N}$ were non-species-specific in the LYCOG experiment. A shift in dominance from a mid-grass to tallgrass species enhanced the ANPP– CO_2 response on the silty clay soil (Fay et al. 2012; Polley et al. 2012b), thereby increasing soil pools of readily decomposable SOM, decomposition rates, and microbial biomass (Procter et al. 2014), and apparently stimulating mineralization of older SOM. Whether responding to an accompanying change in the $\delta^{15}\text{N}$ of plant-available N or to fractionation from other N transformations in soil or plants, $\delta^{15}\text{N}$ values of three prairie species increased by a similar magnitude at elevated CO_2 .

Leaf $\delta^{15}\text{N}$ responded to CO_2 in only one of two 4-year experiments and then only on a silty clay soil on which

leaf $\delta^{15}\text{N}$ values declined at subambient CO_2 and increased slightly at $>450 \mu\text{L L}^{-1} \text{CO}_2$. The CO_2 -caused increase in leaf $\delta^{15}\text{N}$ on the silty clay soil could imply a lesser decline or even an increase in losses of ^{15}N -depleted molecules to leaching or denitrification at elevated CO_2 (Garten et al. 2011), but this interpretation is not compatible with the large and consistently positive ANPP– CO_2 response on the silty clay soil (Polley et al. 2012b) nor with the absence of CO_2 effects on leaf [N]. Alternatively or in addition, CO_2 may have increased foliar $\delta^{15}\text{N}$ on the silty clay soil by reducing fractionation during plant uptake, assimilation, and turnover/export of N (Evans 2001). Change in plant fractionation alone seems an unlikely explanation for $\delta^{15}\text{N}$ trends. In order to have explained results, fractionation must have responded similarly to CO_2 in a forb and two grasses on the silty clay, but not responded to CO_2 on clay and sandy loam soils. Rather, we suggest that CO_2 increased leaf $\delta^{15}\text{N}$ values on the silty clay soil by favoring uptake of N that differed in isotopic signature. Uptake of N differing in isotopic signature may have occurred because CO_2 (1) favored N acquisition from deeper versus shallower soil layers, or (2) changed the $\delta^{15}\text{N}$ signature of source N by increasing microbial immobilization (Billings et al. 2002; Dijkstra et al. 2010) or mineralization of recalcitrant SOM (Gill et al. 2002; Fig. 7). Soil $\delta^{15}\text{N}$ increased with depth in the pasture that we studied, a trend observed in other ecosystems (e.g., Nadelhoffer and Fry 1988). Microbes discriminate against the heavier ^{15}N isotope (Robinson 2001). Microbial fractionation thus enriches the N in recalcitrant forms of SOM (Tiessen et al. 1984). In the LYCOG experiment, CO_2 enrichment increased pools of coarse POM-C by four-fold in the two clay soils, but reduced pools of mineral-associated organic C, generally considered to be recalcitrant to decomposition, on the silty clay soil only (Procter et al., submitted). CO_2 thus may have increased leaf $\delta^{15}\text{N}$ values by increasing mineralization of older, ^{15}N -enriched SOM. Such reallocation of N from soil to plants is consistent with one prediction of the PNL hypothesis (Luo et al. 2004), that CO_2 -caused N limitation may initially be overcome by the transfer of N from SOM with low C:N ratio to plants with higher C:N ratio (Gill et al. 2006). CO_2 enrichment increased $\delta^{15}\text{N}$ values in desert plants (Billings et al. 2002) and the dominant grass of tallgrass prairie (Williams et al. 2006). Billings et al. (2002) attributed greater $\delta^{15}\text{N}$ of desert plants to greater fractionation in soil as a result of increased soil microbial activity at elevated CO_2 . Johnson DW et al. (2000) attributed an increase in foliar $\delta^{15}\text{N}$ of ponderosa pine (*Pinus ponderosa* Dougl.) to greater uptake of N derived from recalcitrant SOM at elevated CO_2 .

Tightening the N cycle could reduce plant $\delta^{15}\text{N}$ by reducing fractionation during N uptake and assimilation by plants (Evans 2001), increasing plant reliance on mycorrhizae for N uptake (Bassirirad et al. 2003; Hobbie and

Hobbie 2008), decreasing the recycling of N through fractionating processes in the soil/plant system, or reducing losses of ^{15}N -depleted N molecules to leaching or denitrification (Garten et al. 2011). Indeed, foliar $\delta^{15}\text{N}$ values declined over the 4 years of both the PCG and LYCOG experiments. This decline in $\delta^{15}\text{N}$ is consistent with the view that fractionation resulting from ecosystem N losses, plant assimilation of N, or other processes declined, or plants exhibited increased reliance on mycorrhizal fungi during community development, regardless of CO_2 treatment (Evans 2001).

On the other hand, our finding that CO_2 did not consistently affect plant $\delta^{15}\text{N}$ is seemingly at variance with results from some CO_2 experiments (Billings et al. 2002; Bassirirad et al. 2003; Garten et al. 2011) and studies in which CO_2 effects on N cycling have been inferred from decadal changes in plant $\delta^{15}\text{N}$ values (Peñuelas and Estiarte 1997; McLauchlan et al. 2010). The absence of a consistent $\delta^{15}\text{N}$ – CO_2 response could imply either that N strongly limited plant growth across experiments and soils or that N cycling and plant N assimilation were relatively insensitive to CO_2 despite apparent tightening of the N cycle during each 4-year experiment. Fractionation of soil N and discrimination during the assimilation of inorganic to organic N in plants are reduced when N is strongly limiting and plant demand approaches N supply rates (Robinson 2001; Billings et al. 2002; Kolb and Evans 2003; Kalcsits et al. 2014). Losses of ^{15}N -depleted N molecules should also be minimal in strongly N-limited systems. Our data on CO_2 effects, however, are most consistent with the interpretation that N cycling and plant fractionation were relatively insensitive to CO_2 during the first 4 years of treatment. Resin-available soil N (NH_4^+ , NO_3^-) decreased by only about 15 % from 280 to 480 $\mu\text{L L}^{-1}\text{CO}_2$ in the LYCOG experiment and was not correlated with ANPP on any soil type (Fay et al. 2012). Similarly, CO_2 did not consistently affect the [N] in aboveground tissues of dominant species from the LYCOG experiment (Polley et al. 2011) and had inconsistent effects on the [N] of dominants from the PCG experiment (Polley et al. 2003). The changes in grassland $\delta^{15}\text{N}$ that did occur along the CO_2 gradient involved changes in species relative abundances, highlighting the underappreciated role of community change in CO_2 –ecosystem interactions.

Author contribution statement All authors contributed to idea formulation, sampling design, and data collection. HWP analyzed data and wrote the manuscript; other authors provided editorial advice.

Acknowledgments Chris Kolodziejczyk operated CO_2 chambers. Field and laboratory assistance from Anne Gibson, Katherine Jones,

Chris Kolodziejczyk, Alicia Naranjo, Kyle Tiner, and numerous students was invaluable. Joseph Craine and Kevin Mueller provided insightful suggestions that improved the manuscript. R.J. acknowledges support from the U.S. Department of Energy (Program in Ecosystem Research #ER64242). Mention of trade names or commercial products does not imply endorsement by the US Department of Agriculture. USDA is an equal opportunity provider and employer.

References

- Bassirirad H, Constable JVH, Lussenhop J, Kimball BA, Norby RJ, Oechel WC, Reich PB, Schlesinger WH, Zitzer S, Sehtiya HL, Silim S (2003) Widespread foliage $\delta^{15}\text{N}$ depletion under elevated CO_2 : inferences for the nitrogen cycle. *Glob Change Biol* 9:1582–1590
- Bechtold JC, Naiman RJ (2006) Soil texture and nitrogen mineralization potential across a riparian toposequence in a semi-arid savanna. *Soil Biol Biochem* 38:1325–1333
- Billings SA, Schaeffer SM, Zitzer S, Charlet T, Smith SD, Evans RD (2002) Alterations of nitrogen dynamics under elevated carbon dioxide in an intact Mojave Desert ecosystem: evidence from nitrogen-15 natural abundance. *Oecologia* 131:463–467
- Cambardella C, Elliott E (1992) Particulate soil organic-matter changes across a grassland cultivation sequence. *Soil Sci Soc Am J* 56:777–783
- Craine JM, Elmore AJ, Aida MPM, Bustamante M, Dawson TE, Hobbie EA, Kahmen A, Mack MC, McLauchlan KK, Michelsen A, Nardoto GB, Pardo LH, Peñuelas J, Reich PB, Schuur EAG, Stock WD, Templer PH, Virginia RA, Welker JM, Wright IJ (2009) Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytol* 183:980–992
- De Graaff MA, van Groenigen KJ, Six J, Hungate B, van Kessel C (2006) Interactions between plant growth and soil nutrient cycling under elevated CO_2 : a meta-analysis. *Glob Change Biol* 12:2077–2091
- Dijkstra FA, Blumenthal D, Morgan JA, Pendall E, Carrillo Y, Follett RF (2010) Contrasting effects of elevated CO_2 and warming on nitrogen cycling in a semiarid grassland. *New Phytol* 187:426–437
- Evans RD (2001) Physiological mechanisms influencing plant nitrogen isotope composition. *Trends Plant Sci* 6:121–126
- Fay PA, Kelley AM, Procter AC, Jin VL, Jackson RB, Johnson HB, Polley HW (2009) Primary productivity and water balance of grassland vegetation on three soils in a continuous CO_2 gradient: initial results from the Lysimeter CO_2 Gradient Experiment. *Ecosystems* 12:699–714
- Fay PA, Jin VL, Way DA, Potter KN, Gill RA, Jackson RB, Polley HW (2012) Soil-mediated effects of subambient to increased carbon dioxide on grassland productivity. *Nat Clim Change* 2:742–746
- Garten CT Jr, Iversen CM, Norby RJ (2011) Litterfall ^{15}N abundance indicates declining soil nitrogen availability in a free-air CO_2 enrichment experiment. *Ecology* 92:133–139
- Gill RA (2007) Influence of 90 years of protection from grazing on plant and soil processes in the subalpine of the Wasatch Plateau, USA. *Rangeland Ecol Manag* 60:88–98
- Gill RA, Polley HW, Johnson HB, Anderson LJ, Maherali H, Jackson RB (2002) Nonlinear grassland responses to past and future atmospheric CO_2 . *Nature* 417:279–282
- Gill RA, Anderson LJ, Polley HW, Johnson HB, Jackson RB (2006) Potential nitrogen constraints on soil carbon sequestration under low and elevated atmospheric CO_2 . *Ecology* 87:41–52

- Grace JB (2006) Structural equation modeling and natural systems. Cambridge University Press, Cambridge
- Gu C, Riley WJ (2010) Combined effects of short term rainfall patterns and soil texture on soil nitrogen cycling – a modeling analysis. *J Contam Hydrol* 112:141–154
- Handley LL, Scrimgeour CM (1997) Terrestrial plant ecology and ^{15}N natural abundance: the present limits to interpretation for uncultivated systems with original data from a Scottish old-field. *Adv Ecol Res* 27:133–212
- Hobbie EA, Hobbie JE (2008) Natural abundance of ^{15}N in nitrogen-limited forests and tundra can estimate nitrogen cycling through mycorrhizal fungi: a review. *Ecosystems* 11:815–830
- Hungate BA, Stiling PD, Dijkstra P, Johnson DW, Ketterer ME, Humus GJ, Hinkle CR, Drake BG (2004) CO_2 elicits long-term decline in nitrogen fixation. *Science* 304:1291
- Johnson DW, Cheng W, Ball JT (2000) Effects of CO_2 and N fertilization on decomposition and N immobilization in ponderosa pine litter. *Plant Soil* 229:203–212
- Johnson HB, Polley HW, Whitis RP (2000) Elongated chambers for field studies across atmospheric CO_2 gradients. *Funct Ecol* 14:388–396
- Kahmen A, Wanek W, Buchmann N (2008) Foliar $\delta^{15}\text{N}$ values characterize soil N cycling and reflect nitrate or ammonium preference of plants along a temperate grassland gradient. *Oecologia* 156:861–870
- Kalcsits LA, Buschhaus HA, Guy RD (2014) Nitrogen isotope discrimination as an integrated measure of nitrogen fluxes, assimilation and allocation in plants. *Physiol Plant* 151:293–304
- Kelley AM, Fay PA, Polley HW, Gill RA, Jackson RB (2011) Atmospheric CO_2 and soil extracellular enzyme activity: A meta-analysis and CO_2 gradient experiment. *Ecosphere* 2(8):art96
- King JY, Mosier AR, Morgan JA, LeCain DR, Milchunas DG, Parton WJ (2004) Plant nitrogen dynamics in shortgrass steppe under elevated atmospheric carbon dioxide. *Ecosystems* 7:147–160
- Kolb KJ, Evans RD (2003) Influence of nitrogen source and concentration on nitrogen isotope discrimination in two barley genotypes (*Hordeum vulgare* L.). *Plant Cell Environ* 16:1431–1440
- Littell RS, Stroup WW, Freund RJ (2002) SAS for linear models, 4th edn. SAS Institute, Cary
- Luo Y, Su B, Currie WS, Dukes JS, Finzi A, Hartwig U, Hungate B, McMurtrie RE, Oren R, Parton WJ, Pataki DE, Shaw MR, Zak DR, Field CB (2004) Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *Bioscience* 54:731–739
- McLauchlan KK, Ferguson CJ, Wilson IE, Ocheltree TW, Craine JM (2010) Thirteen decades of foliar isotopes indicate declining nitrogen availability in central North American grasslands. *New Phytol* 187:1135–1145
- Mielnick PS, Dugas WA, Johnson HB, Polley HW, Sanabira J (2001) Net grassland carbon flux over a subambient to superambient CO_2 gradient. *Glob Change Biol* 7:747–754
- Nadelhoffer KJ, Fry B (1988) Controls on natural ^{15}N and ^{13}C abundances in forest soil organic matter. *Soil Sci Soc Am J* 52:1633–1640
- Peñuelas J, Estiarte M (1997) Trends in plant carbon concentration and plant demand for N throughout this century. *Oecologia* 109:69–73
- Polley HW, Johnson HB, Derner JD (2002) Soil- and plant-water dynamics in a C_3/C_4 grassland exposed to a subambient to superambient CO_2 gradient. *Glob Change Biol* 8:1119–1129
- Polley HW, Johnson HB, Derner JD (2003) Increasing CO_2 from subambient to superambient concentrations alters species composition and increases above-ground biomass in a C_3/C_4 grassland. *New Phytol* 160:319–327
- Polley HW, Johnson HB, Fay PA, Sanabria J (2008) Initial response of evapotranspiration from tallgrass prairie vegetation to CO_2 at subambient to elevated concentrations. *Funct Ecol* 22:163–171
- Polley HW, Fay PA, Jin VL, Combs GF Jr (2011) CO_2 enrichment increases element concentrations in grass mixtures by changing species abundances. *Plant Ecol* 212:945–957
- Polley HW, Jin VL, Fay PA (2012a) CO_2 -caused change in plant species composition rivals the shift in vegetation between mid-grass and tallgrass prairies. *Glob Change Biol* 18:700–710
- Polley HW, Jin VL, Fay PA (2012b) Feedback from plant species change amplifies CO_2 enhancement of grassland productivity. *Glob Change Biol* 18:2813–2823
- Polley HW, Derner JD, Jackson RB, Wilsey BJ, Fay PA (2014) Impacts of climate change drivers on C_4 grassland productivity: scaling driver effects through the plant community. *J Exp Bot* 65:3415–3424
- Procter AC, Ellis JC, Fay PA, Polley HW, Jackson RB (2014) Fungal community responses to past and future atmospheric CO_2 differ by soil type. *Appl Environ Microbiol* 80:7364–7377
- Robinson D (2001) $\delta^{15}\text{N}$ as an integrator of the nitrogen cycle. *Trends Ecol Evol* 16:153–162
- Robinson D, Conroy JP (1999) A possible plant-mediated feedback between elevated CO_2 , denitrification and the enhanced greenhouse effect. *Soil Biol Biochem* 31:43–53
- Shipley B (2000) Cause and correlation in biology: a user's guide to path analysis, structural equations and causal inference. Cambridge University Press, Cambridge
- Soper FM, Boutton TW, Sparks JP (2014) Investigating patterns of symbiotic nitrogen fixation during vegetation change from grassland to woodland using fine scale $\delta^{15}\text{N}$ measurements. *Plant Cell Environ*. doi:10.1111/pce.12373
- Stock WD, Evans JR (2006) Effects of water availability, nitrogen supply and atmospheric CO_2 concentrations on plant nitrogen natural abundance values. *Funct Plant Biol* 33:219–227
- Tiessen H, Karamanos RE, Stewart JWB, Selles F (1984) Natural nitrogen-15 abundance as an indicator of soil organic matter transformations in native and cultivated soils. *Soil Sci Soc Am J* 48:312–315
- Wang L, Macko SA (2011) Constrained preferences in nitrogen uptake across plant species and environments. *Plant Cell Environ* 34:525–534
- Williams MA, Rice CW, Owensby CE (2006) Natural ^{15}N abundances in a tallgrass prairie ecosystem exposed to 8-y of elevated atmospheric CO_2 . *Soil Biol Biochem* 28:409–412